

**Notes on the Vertebrates of northern Pará, Brazil:  
a forgotten part of the Guianan Region, I. Herpetofauna**  
Notas sobre os vertebrados do norte do Pará, Brasil:  
uma parte esquecida da Região das Guianas, I. Herpetofauna

Teresa Cristina Sauer Avila-Pires<sup>I</sup>  
Marinus Steven Hoogmoed<sup>II</sup>  
Wáldima Alves da Rocha<sup>III</sup>

**Abstract:** We discuss the herpetological results of seven expeditions to the Guianan part of Pará, which resulted in a total of 80 species of amphibians (77 frogs and three caecilians) and 95 species of reptiles (36 species of lizards, three species of amphisbaenians, 49 species of snakes, five species of chelonians and two species of caiman). We report six species new to science (three frogs, one caecilian, one lizard, one amphisbaenian), six new records for Brazil (five frogs, one caecilian) and 23 new records for Pará (13 frogs, four lizards, six snakes). For each of the new records we provide comments. Special comment is made about a large population of the toad *Atelopus hoogmoedi* that seems to be doing well and does not show any signs of population decline as many species of *Atelopus* at higher elevations do. We provide a complete list of species collected per locality containing data on endemism, habitat, reproduction and food. For each of the seven collecting sites we provide data on richness and abundance of species. The sites are compared regarding their species composition, even though we can not say how much of the differences are due to specific habitats or geographic variation, seasonal variation or sampling deficiency. We synonymised the Bufonid *Rhinella martyi* with *Bufo margaritifera* and selected a lectotype for *Rana margaritifera* in order to resolve the problems about this name.

**Keywords:** Amphibia. Reptilia. Guiana Centre of Endemism. Brazil. Species richness. Species list.

**Resumo:** Os resultados herpetológicos de sete expedições à parte guianense do estado do Pará são apresentados e discutidos, registrando-se um total de 80 espécies de anfíbios (77 anuros e três Gymnophiona) e 95 espécies de répteis (36 espécies de lagartos, três espécies de anfisbênídeos, 49 espécies de ofídios, cinco espécies de quelônios e duas espécies de jacarés). Dessas espécies, seis são novas para a ciência (três anuros, um Gymnophiona, um lagarto, um anfisbênídeo), seis representam novos registros para o Brasil (cinco anuros, um Gymnophiona) e 23 novos registros para o Pará (13 anuros, quatro lagartos, seis ofídios). Comenta-se cada um dos novos registros. Comentários especiais são feitos sobre uma grande população do sapo *Atelopus hoogmoedi*, a qual parece estar bem saudável e não mostra sinais de declínio populacional, como muitas espécies de *Atelopus* em outros lugares de maior altitude. Uma lista completa das espécies coletadas por localidade, incluindo dados sobre endemismo, habitat, reprodução e alimentação, é apresentada. Para cada uma das sete áreas de coleta, apresentamos dados sobre riqueza e abundância de espécies. As áreas são comparadas quanto à similaridade na composição das espécies, ainda que não seja possível indicar quanto das diferenças encontradas deve-se a ambientes específicos ou variação geográfica, variação sazonal ou deficiência na amostragem das espécies. O Bufonidae *Rhinella martyi* é considerado sinônimo de *Bufo margaritifera* e um lectótipo para *Rana margaritifera* é selecionado visando dirimir dúvidas sobre o nome da espécie.

**Palavras-chave:** Amphibia. Reptilia. Centro de Endemismo Guiana. Brasil. Riqueza de espécies. Lista de espécies.

---

<sup>I</sup> Museu Paraense Emílio Goeldi. Belém, Pará, Brasil (avilapires@museu-goeldi.br).

<sup>II</sup> Museu Paraense Emílio Goeldi. Belém, Pará, Brasil (marinus@museu-goeldi.br).

<sup>III</sup> Universidade Aberta do Brasil. Canto do Buriti, Piauí, Brasil (waldima@yahoo.com.br).



## INTRODUCTION

On December 4, 2006, the State of Pará created five new conservation units in the northern part of Pará, north of the Amazon, in order to establish a large and protected, mostly forested, area that would form a continuous block with similarly protected areas in Amapá (Parque Nacional [PARNA] Montanhas de Tumucumaque), French Guiana, Suriname and Guyana, and with Indian Territories in the region (Figure 1) (Governo do Estado do Pará, 2006). The five conservation units created by the state were: Estação Ecológica (ESEC) Grão-Pará (4.2 million ha), Reserva Biológica (REBIO) Maicuru (1.2 million ha), and the Florestas Estaduais (FLOTA) de Faro (0.6 million ha), Trombetas (3.2 million ha) and Paru (3.6 million ha). Together they cover an area of 13.2 million ha and, with the already existing protected areas – Indian territories (TI) of Trombetas-Mapuera, Tumucumaque, rio Paru d'Este, Nhamundá-Mapuera and Zo'é; two 'Quilombola' (African-Brazilian) territories; the Florestas Nacionais (FLONA) Saracá-Taquera and da Mulata, REBIO do rio Trombetas, and ESEC Jari – they form an enormous block, although with different degrees of protection (besides TI and 'quilombola' sites, that harbour traditional populations, FLOTA's and FLONA's aim at the sustainable use of natural resources). Most of the newly created conservation units are covered by non-flooded tropical rainforest ('terra-firme' forest), but in several places other vegetation types, like flooded forests ('várzea' and 'igapó'), savanna and 'cerradão', are present as well. Only a relatively narrow band of land in 'Calha Norte Paraense' (CNP) along the Amazon is not protected and open to unregulated human occupation. We here consider as CNP that part of Pará that is situated north of the Amazon River.

On February 14, 2007 several parties (Government of Pará [SEMA-PA], Conservação Internacional [CI-Brasil], Museu Paraense Emílio Goeldi [MPEG], Instituto do Homem e Meio Ambiente da Amazônia [IMAZON], Instituto de Desenvolvimento Florestal do Estado do Pará [IDEFLOR], Instituto de Manejo e Certificação Florestal e Agrícola

[IMAFLORA], and the German Technical Cooperation Agency [GTZ]) signed an agreement to form a consortium to provide data to develop management plans for the five state protected areas ('Projeto Diagnóstico da Biodiversidade das Unidades de Conservação Estaduais do Mosaico Calha Norte, Estado do Pará'). As a result of this agreement seven expeditions to investigate the biodiversity of the recently established protected areas were planned. The localities to be inventoried were chosen by specialists from SEMA-PA, CI-Brasil, IMAZON and MPEG, on the basis of satellite photographs, georeferenced databases, and vegetation and altitude data, with the goal to optimize the coverage of the different phytophysionomies and altitudes present in the area during seven three-week long expeditions distributed over a year. An additional overflight helped with the final definition of the areas chosen for sampling. As final transportation to five of the research areas was planned by helicopter, the localities could be chosen independent of road or river transport. Fieldwork started in January 2008 and ended in January 2009. During this period three expeditions to different localities in ESEC Grão-Pará and one each to a locality in the other four newly created conservation units were carried out by 12 – 13 researchers of the MPEG and about ten technical assistants.

Due to its continental size, many Amazonian areas are still poorly known regarding their herpetofauna. The northern part of the state of Pará, in Brazil, is one of these areas, with only a few spots reasonably well surveyed. This area north of the Amazon River forms part of the Guianan Region (or 'Guianas') as defined by Hoogmoed (1979b), mainly delimited by the Orinoco, Negro and Amazonas rivers on the west and south, and the Atlantic Ocean on the north and east. The Guianan Region encompasses the three Guianas (Guyana, Suriname, French Guiana), the southeastern part of Venezuela, and in Brazil the states of Amapá and Roraima, the state of Pará north of the Amazon River, and the relatively small northeastern part of the state of Amazonas, north of the Amazon River and east and north of the rio Negro. Silva *et al.* (2005) considered the area as



the Guianan Area of Endemism, the largest of eight areas of endemism in the Amazon region, with half of its surface in Brazil. According to Silva *et al.*, (2005) in the Brazilian part of this area of endemism, only 4.06% had been deforested, although this percentage may have gone up in the past few years. We do not include part of Colombia in the Guianan Region (and neither do Silva *et al.*, 2005), as was done by the 2002 Paramaribo workshop (Hollowell & Reynolds, 2005 and articles therein), as there are no good zoogeographical reasons for that inclusion. Concerning herpetofauna, the area west of the rio Negro has no Guiana endemics. The Guianan Region has a number of species in common with other areas of Amazonia, but also has a number of endemic species. This is especially true for the 'tepui', sandstone mountains with elevations above 1,500 m, usually considered a distinct biogeographic region ('Pantepui') within the Guianan Region (Hoogmoed, 1979b; McDiarmid & Donnelly, 2005), in Venezuela, western Guyana and extreme northern Brazil (Roraima and Amazonas States). In Suriname there is one sandstone mountain of 1,200 m altitude, but it has no Pantepui endemics (MSH, pers. obs.), which generally only occur above 1,500 m. In northern Pará, Amapá and French Guiana no tepuis are found and elevations just reach 900 m, thus explaining the absence of any herpetological tepui endemics in the area. A number of lowland species are also endemic to the Guianan region or part of it, although numbers have dropped when new range extensions became available (e.g. Caldwell & Hoogmoed, 1998).

About 350 species of amphibians and a similar number of reptiles are known from the whole of Amazonia, including the Guianas (see Eva & Huber, 2005: 11 for the limits of Amazonia, here considered as the area named Amazonia *sensu lato* [Ia+IIa+IIb]), of which c. 82% of the amphibians and 62% of the reptiles are endemic (Avila-Pires *et al.*, 2007; Duellman, 1999). Hoogmoed (1979b) estimated that, for the Guianan lowlands, c. 52% of the amphibians and 26% of the reptiles were endemic. However, as more areas were better surveyed,

many of the lowland species considered endemic to the Guianas in 1979 were shown to have a wider distribution throughout Amazonia (e.g. *Allophryne ruthveni* [Caldwell & Hoogmoed, 1998], *Bufo guttatus* and *Lithodytes lineatus* [MSH, unpublished data, material in MPEG]). Including also the fauna of the tepuis, Señaris & MacCulloch (2005) found that 54% of the amphibian species from the Guianas were endemic to the region, while Avila-Pires (2005) indicated that 30% of the reptile species were endemic. As there are still large gaps in our knowledge about the herpetofauna of the Guianan Region, both range extensions and new species are expected to be found in northern Pará.

A number of recent herpetofaunal studies focus on (part of) the Guianan Region, among them those on amphibians (Fouquet *et al.*, 2007a, b; Lescure & Marty, 2000; Kok, 2000; Kok *et al.*, 2006a), lizards (Hoogmoed & Lescure, 1975; Hoogmoed & Avila-Pires, 1989; Gasc, 1990), amphisbaenians and snakes (Gasc & Rodrigues, 1980; Chippeaux, 1986; Starace, 1998) from French Guiana; anurans (Hoogmoed, 1969a, b, 1971a, b, 1979a), wormsalamanders (Nussbaum & Hoogmoed, 1979), lizards and amphisbaenians (Hoogmoed, 1973), and some groups of snakes (Hoogmoed, 1977, 1980, 1983, 1985) from Suriname; an increasing number of studies on the herpetofauna of Guyana, especially the western (Pantepui) part of this country (Cole & Kok, 2006; Kok, 2005, 2006a, b, 2008a, b, 2009; Kok *et al.*, 2006b; Kok & Castroviejo-Fisher, 2008; Kok & Kalamandeen, 2008; Lathrop & MacCulloch, 2007; MacCulloch & Lathrop, 2001, 2002, 2004a, b, 2005, 2009; MacCulloch *et al.*, 2006, 2007, 2008a, b), several from the central and southern Guyana lowlands, like the Mabura Hill and Iwokrama region (Donnelly *et al.*, 1998, 2005a, b, 2006; Ernst *et al.*, 2005, 2007; Kok & Ernst, 2007, Kok *et al.*, 2007, Señaris *et al.*, 2008) and from the coastal area (EMC, 2006); the study by Gorzula & Señaris (1999) on the herpetofauna of Venezuelan Guayana, and by Pritchard & Trebbau (1984) on chelonians from Venezuela; and those on amphibians (Lima *et al.*, 2006), snakes (Martins



& Oliveira, 1993, 1998), and lizards (Vitt *et al.*, 2008) from the Manaus area, Amazonas, Brazil. More specific papers are those by Hoogmoed & Avila-Pires (1991a), with data on Amphisbaenidae; by Hoogmoed & Avila-Pires (1992) on the lizard genus *Arthrosaura*, by Cunha *et al.* (1980), Carvalho (1997, 2002) and Vanzolini & Carvalho (1991), on lizards and snakes from Roraima. Some publications deal with species that occur throughout a large part of the Guianan Region (Campbell & Lamar, 2004; Dixon *et al.*, 1993; Medem, 1983; Noonan & Gaucher, 2005, 2006; Noonan & Wray, 2006; Roze, 1996; Wollenberg *et al.*, 2006, 2008). Conservação Internacional organized a series of expeditions to the Tumucumaque Mountains on the border of French Guiana and Amapá, between 2004 and 2006, but the herpetological results (Lima, 2008) are still under discussion and at the moment only can be used with much care, checking each species record. Hoogmoed (1979b, 1983), Hoogmoed & Avila-Pires (1991b), Señaris & MacCulloch (2005) and Avila-Pires (2005) present lists of Guianan herpetofauna, including data from Amapá, Amazonas, Pará and Roraima. In Avila-Pires (1995), a catalogue of the lizards of Brazilian Amazonia, data on lizards from the Guianan Region can also be found, while data on chelonians may be found in Vogt (2008). Bartlett & Bartlett (2003) is a good general introductory book for the Amazonian herpetofauna, and Rueda-Almonacid *et al.* (2007) is an excellent fieldguide for chelonians and crocodiles, but both are not complete for the Guianan Region. Avila-Pires *et al.* (2009) compared the lizard faunas from three sites on the Guiana Shield (Brokopondo and Sipaliwini [both Suriname, the last one on the border with Brazil] and Balbina [Amazonas, Brazil]) with ten sites in other areas of the Amazon Region. They showed that the Guianan sites were most closely related to Belém and Caxiuana (both in Pará) in the lower Amazon area.

A list of species of amphibians and reptiles present in CNP as a whole does not exist. The areas that have been better studied are the lower Trombetas River, where environmental studies have been done in the context of a

large bauxite mining project in the area (U. Galatti, pers. obs.); and the Jari (Monte Dourado) area, at the border with Amapá state, as part of a two-year multidisciplinary study by C. Peres and T. A. Gardner of the University of East Anglia (United Kingdom), and collaborators. Most data on the Trombetas studies, however, are not published. Part of the results of the Jari project, regarding herpetofauna, can be found in Gardner *et al.* (2007) and Ribeiro-Junior *et al.* (2008). Hoogmoed and Avila-Pires made collections (material in Museu Paraense Emílio Goeldi, Belém, Pará [MPEG] and in the National Museum of Natural History, Leiden, The Netherlands [RMNH]) in the rio Nhamundá area in 1988, the results of which have only partly (lizards) been published by Avila-Pires (1995). Between 1980 and 2006 personnel of MPEG made several collecting trips to the municipalities Almeirim and Monte Alegre and obtained small, but interesting collections, which are now in MPEG. Besides, occasional expeditions have been made, especially following the large rivers, which account for the sparse data found in the literature and specimens in collections. Avila-Pires (1995) registered a number of lizards from this area, even though only from few localities, showing large gaps of information for the area as a whole. Vogt (1994, 2008) and Haller & Rodrigues (2005, 2006) give data on chelonian species from the Trombetas River.

Based on the existing literature, for CNP we may expect approximately 100 species of anurans and up to nine species of Gymnophiona (Lescure & Marty, 2000; Lima *et al.*, 2006; Señaris & MacCulloch, 2005; A.O. Maciel & Hoogmoed, unpublished data). Among reptiles, we could expect about 40 species of lizards, ten species of amphisbaenians, 100 of snakes, 11 of chelonians, and three species of caimans (Hoogmoed, 1973; Chippeaux, 1986; Martins & Oliveira, 1993, 1998; Avila-Pires, 1995, 2005; Starace, 1998; Rueda-Almonacid *et al.*, 2007; Vogt, 2008; MSH, unpublished data). As no tepuis are present in the CNP area, tepui endemics are not expected to be found there, only lowland species (< 750 m). On the other hand, a number of species not present in other Guianan countries (French Guiana, Suriname, Guyana and Venezuela)



possibly can be expected to occur in the areas under influence of the Amazon River.

The study here presented as part of the CNP Project intended to inventory the herpetofauna from key localities surveyed during seven expeditions (Figure 2), taking into account the necessity to produce management plans for the five conservation units created by the State of Pará, Brazil, in 2006 – Floresta Estadual de Faro (FLOTA Faro), Floresta Estadual do Paru (FLOTA Paru), Floresta Estadual do Trombetas (FLOTA Trombetas), Reserva Biológica de Maicuru (REBIO Maicuru) and Estação Ecológica do Grão-Pará (ESEC Grão-Pará). Considering the large extension of the area covered by the five conservation units (13.2 million ha), it was impossible, within the period of 13 months, to accomplish intensive studies on the fauna of each of them. Therefore it was decided to select a number of points that together could cover the different phytophysionomies encountered in northern Pará, as explained before, and to perform in each of them a Rapid Assessment Program (RAP). Although the results obtained are not exhaustive, and new studies will be necessary to improve our understanding of the herpetofauna of the area, they represent an important advance in our knowledge, and provide the basis for management plans. We present here an analysis based on all expeditions, because much information is common to all or is complementary, and after that we highlight several species that represent new or interesting zoogeographic data.

## MATERIAL AND METHODS

### STUDY AREAS

The greater part of CNP is on the Guiana Shield, only a wide band north of the Amazon belongs to the alluvial Amazon valley. The core of the Guiana Shield is made up of pre-Cambrian metamorphic and igneous rocks, especially granites and gneisses. On all sides the core of the Guiana Shield is surrounded by a band of low areas of varying width consisting of alluvial sediments. The higher

part of the Guiana Shield is covered with sandstone remnants of the Roraima Formation, which was deposited in Proterozoic time, 1.6-1.8 billion years ago. After uplift, this formation covered the Guiana Shield as an extensive sandstone plateau or tableland. During the Late Cretaceous and in the Tertiary there were new periods of further uplift of the area, at the same time that erosion shaped the present-day table mountains or tepuis, which are concentrated in the NW part of the Guiana Shield, in SE Venezuela and adjacent W Guyana, with some tepuis on the border of Brazil (Roraima and Amazonas States) with these countries (Hoogmoed, 1979b and literature cited therein). In northern Pará no sandstone tepuis are present, and consequently, by definition, no herpetofaunal tepui endemics. The northern and southern part of the Guianan Region are separated by the divide between rivers that are part of the Amazon basin and flow S from the divide to that river, and rivers of the Guianas that flow north directly to the Atlantic Ocean. The mountains of the divide, which is formed by the Acarai Mountains in the West (between Brazil and Guyana) and the Tumucumaque Mountains in the East (between Suriname and French Guiana on one side and Brazil on the other) are relatively low (in some places, like the Sipaliwini area, not higher than 250 m, in one place reaching up to about 900 m, but generally below 800 m). From the Amazon River the area of northern Pará gradually slopes north, up towards the divide. The area is hilly, with rounded hills and elevated plateaus at a level of about 500 m, that at least in part contain bauxite deposits. All rivers in the area run roughly N-S and have numerous rapids and waterfalls from their upper reaches to close to the Amazon, and therefore are difficult to navigate. The Trombetas River in the South and the Suriname River in the north (Hoogmoed, 1973) seem to divide the Guianan Region in an eastern and a western part and may form a distribution barrier for some species.

Most areas we inventoried have a wet tropical climate (Am according to the Köppen classification), but site ESEC Grão-Pará North is in an area that is



characterized as Aw (Peel *et al.*, 2007; SUDAM, 1984). The rainy season generally is between December and June, with a short drier break in February/March, the dry season is between July and November. Total mean annual rainfall is around 2,100-2,500 mm per year for most areas inventoried, except for sites ESEC Grão-Pará North and ESEC Grão-Pará Centre, which are in areas with 2,000-2,100 mm, and site FLOTA Paru, which is in an area with a mean annual rainfall of 1,500-2,000 mm (Figure 3). Mean annual temperatures in most of the area are about 25-26 °C, but sites FLOTA's Faro, Trombetas and Paru are in areas where that temperature is about 27 °C (Figure 4) (SUDAM, 1984).

The study areas are located in the Guianan Region of the northern part of Pará, north of the Amazon River, in the five conservation units established in December 2006: one each in the FLOTA's Faro, Trombetas and Paru, one in the REBIO Maicuru and three in the ESEC Grão-Pará (Figure 2). Because of its large size, covering several vegetation types, three expeditions were made to ESEC Grão-Pará: one to the most northwestern part, close to the frontier with Guyana, one to the central part, just south of the Indian Territory of Tumucumaque, and one to the southeastern part, close to the border with FLOTA Paru. A short description of the research areas is provided below. Unfortunately, no details on the vegetation have been provided by the botanists yet, so general terms are used.

FLOTA Faro (0.6 million ha) is situated in the municipalities of Faro and Oriximiná, on the right bank of the rio Nhamundá, which forms the border of Pará with Amazonas. The area belongs to the rio Nhamundá basin and is covered by tropical rain forest (terra-firme forest = 'Floresta Ombrófila Densa das Terras Baixas' according to RADAM-Brasil ["Radar na Amazônia" project] terminology), except in a band of about 700 m along the river and some distance up along the creeks, that is covered by várzea forest ('Floresta Ombrófila Densa Aluvial'), which is inundated during part of the year (Figure 5). Four trails (all in the municipality of Faro) were cut, radiating from a base

camp (S 1° 42' 50.44" W 57° 12' 47.88") that was located on the northern (left) bank of the rio Nhamundá, some distance WNW of Faro, where the river runs more or less east-west. Trails 1-3 ran in a northerly direction, parallel to each other and separated by 800 m, for a distance of 3 km. They started out in várzea forest and after about 700 m entered terra-firme forest. Trail 4 ran 1.5 km SW and 1.5 km SE of the camp, following the riverbank through várzea forest. The area studied was between 0 and 30 m above sea level, with low hills. The river is in open contact with the Amazon, no rapids or waterfalls being present downstream from the collecting area. Pitfall traps were installed in each of the trails 1, 2 and 3, with a distance of 250 m between them within each trail. Because of inundation, no pitfalls were installed in trail 4. The first two pitfalls in trail 1 were inundated and did not work. Trail 1 had pitfalls at 50 m, 250 m, 500 m and 750 m. Trail 2 had pitfalls at 600 m, 850 m, 1,100 m and 1,350 m. Trail 3 had pitfalls at 200 m, 450 m, 700 m, 950 m, 1,200 m, 1,450 m, 1,700 m and 1,950 m. FLOTA Faro was sampled during the first expedition, between January 14 and 28, 2008, during the wet season. At that time the level of the river was high.

FLOTA Trombetas (3.2 million ha) is situated in the municipalities of Oriximiná, Óbidos and Alenquer. It forms part of the basin of the Trombetas River and in the north it borders on the western part of ESEC Grão-Pará. Our camp (S 0° 57' 45.97" W 55° 31' 20.28") was in the municipality of Óbidos, in the southeastern corner of the unit, in an area mainly covered by terra-firme forest ('Floresta Ombrófila Aberta Submontana'). The camp was situated in an opening in a forest with many old *Cecropia* trees, at about 100 m from a creek with rocks and a sandy bottom. It was at the base of a higher area with large rocks and an open forest resembling secondary forest ('capoeira'). Three trails radiated from the camp. Trail 1 ran initially E for 2 km, passing the helicopter landing area at 500 m and then turned N for 7 km, reaching the rio Cuminapanema in a transitional area. Trail 2 ran W for 5 km, first rather level, but after crossing two small creeks steadily uphill, in the last



few hundred meters reaching 'Floresta Ombrofila Densa Submontana'. Trail 3 ran SE for 3 km, the first kilometer up a hill (the same one with the helicopter landing area on top) that was strewn with large granite boulders and covered by a low, open type of forest with lianas (Figure 6), and then descending into a lower, flat area with several creeks. The area studied was between 300 and 450 m above sea level, and was hilly, with a number of small, shallow, clear water creeks, sometimes with steep banks, cut about 10 m into the surrounding terrain. The helicopter landing area was at an altitude of 350 m on a rocky hill top with rockslates (lajedos) and an open vegetation of low bushes, cactus (*Cereus*) and bromeliads (Figure 7). This area formed a distinctive habitat, quite different from the surrounding terra-firme forest. Some large rockslates and boulders also were present in part of the adjacent terra-firme forest (between the helicopter landing area and the camp), which caused some of the open habitat and rockdwelling species to enter the forest. Pitfalls and driftfences were placed in trail 1 at 100 m, 400 m and 950 m from camp. In trail 2 at 200 m, 450 m, 700 m, 1,000 m, 1,250 m, 1,500 m, 1,750 m, 2,100 m, 3,250 m, 3,500 m, 3,750 m, 4,000 m and 4,250 m. The pitfalls at distances over 3,250 m on April 19, 2008 were relocated to 300 m, 1,100 m, 1,350 m, 1,600 m and 1,900 m because of logistical problems with pitfalls beyond 2,100 m. In trail 3 no pitfalls were placed. FLOTA Trombetas was the target of the second expedition, which took place between April 16 and May 1, 2008, in the middle of the rainy season.

REBIO Maicuru (1.2 million ha) is situated in the municipalities of Almeirim and Monte Alegre and is drained by the rivers Maicuru, Paru and Jari. On the north-northeast it reaches the rio Jari (border with Amapá), on the southeast and south it is bordered by FLOTA Paru, and on the west (and partly northwest) it borders on the ESEC Grão-Pará (for a short distance), and the Indian Territories (TI) Rio Paru d'Este and Tumucumaque. Our camp (N 0° 49' 43.03" W 53° 55' 52.32") was located in the municipality of Almeirim, in the middle of the conservation unit, at an

altitude of 150 m above sea level, at some distance from the rio Ipitinga and about 15 m above the river plain. The research area was covered with terra-firme forest ('Floresta Ómbrofila Densa Submontana') (Figure 8), but the forest along the river apparently was regularly flooded, as shown by high water marks on the vegetation. This river forest (igapó) differed from terra-firme forest by being denser, with many low-slung lianas and growth of smaller trees and in some places by the presence of large *Guadua* bamboo stands. The banks of the river were steep, but in several places there were sandy beaches that dropped steeply in the water (Figure 9). From the camp, trail 1 ran NNW for 4.7 km, parallel to, but at some distance from, the river, through terra-firme forest; at about 1 km from camp it crossed a large inundated area along a creek. Trail 2 ran WNW for 5 km through terra-firme forest in terrain with steep hills and ridges. Trail 3 ran SW for 6 km, steadily climbing and near its end reached an altitude of about 550 m. Trail 4 ran S, closely following the river bank through regularly flooded (dry at the time) forest. The area was rather flat, with isolated small hills, but in the SW part there was a large hill-complex reaching an altitude of 550 m. Pitfalls and driftfences were placed in trail 1 at 360 m, 600 m, 800 m and 950 m from camp. In trail 2 they were placed at 350 m, 500 m, 700 m and 900 m from camp. Both trails 1 and 2 ran through terra-firme forest. In trail 3, also through terra-firme forest, no pitfalls were installed. Trail 4 ran through regularly flooded (dry at the time) forest close to the riverbank. In order to get a comparable effort in terra-firme and river bank forest, eight pitfalls were installed in trail 4, at 370 m, 750 m, 900 m, 1,040 m, 1,450 m, 1,600 m, 1,800 m and 2,000 m from camp. The fifth expedition visited REBIO Maicuru between October 21 and November 6, 2008, during the dry season.

FLOTA Paru (3.6 million ha) is situated in the municipalities of Monte Alegre, Alenquer and Óbidos and is drained by the rivers Maicuru, Paru and Jari. On the north it is bordered by the eastern part of ESEC Grão-Pará and REBIO Maicuru, on the east it is bordered by the rio Jari,



that forms the border with Amapá. Our camp (S 0° 56' 38.29" W 53° 14' 10.68") was located in the municipality of Almeirim, in the SE part of the FLOTA, where its border is formed by the rio Paru (Figure 10). The camp was situated at about 100 m from the W bank of the river, in a large patch of low secondary vegetation, and on the S and E side it was bordered by a belt of dense liana forest of 500 m wide. Along the river there was an irregular band of 'Floresta Ombrófila Aberta de Terras Baixas' with antropogenic influences, in several places caused by the presence of isolated houses on this bank of the river. The vegetation on the riverbank itself was rather open, apparently regularly flooded ('igapó'), with hardly any undergrowth in the forest, although in some places there were large clearings, completely taken over by grass and bamboo. The banks of the river were gently sloping and the water of the river was clear, but not very transparent. Some distance downriver from our camp there was a complex of rapids, separating this part of the river from direct contact with the waters of the Amazon. The vegetation away from the river consisted of terra-firme forest ('Floresta Ombrófila Densa de Terras Baixas'). The area along the riverbank was flat, but at about one kilometer from the river bank became hilly with steep slopes, no plateaus, and traversed by several large and small creeks with clear, transparent water. From the camp, trail 1 ran for 5 km NE through 'Floresta Ombrófila Aberta de Terra Baixa', crossing one creek and ending on the river bank upstream from the camp. There were many signs of human activities in this area, with hunting trails, felled trees, open areas and overgrown agricultural fields. Trail 2 ran NW for 9 km through terra-firme forest, crossing several creeks and with many changes in altitude. Trail 3 ran SW for 4 km, first crossing the liana forest around the camp, after 1 km it reached an open cultivated area and then ran through terra-firme forest, crossing a partly dry creek with isolated pools of water in a rocky bed and reaching a creek with running water. In the creeks there were rock outcrops. Trail 4 branched off from trail 3 at the open cultivated area and then ran S for 4 km, closely following the river bank

through open river-bank vegetation ('Floresta Ombrófila Aberta de Terra Baixa'), and crossing some creeks. Altitude in the study area varied from 30 to 100 m. Four pitfalls were placed along each trail at distances of 250 m, 500 m, 750 m and 1,000 m, all in level terrain of about 30 – 90 m. The sixth expedition inventoried FLOTA Paru between December 4 and 19, 2008, at the beginning of the rainy season.

ESEC Grão-Pará occupies 4.2 million ha in the municipalities Oriximiná, Óbidos, Alenquer and Monte Alegre, and runs from the frontier with Guyana in the NW to the TI Tumucumaque, TI Rio Paru d'Este and REBIO Maicuru in the E. A large, more or less triangular western part connects by a narrow neck to a smaller more or less rectangular eastern part. On the south it is bordered by the TI Trombetas-Mapuera, FLOTA Trombetas, TI Zo'é and FLOTA Paru. Because of its great size and different physionomies three localities were sampled in this conservation unit, respectively ESEC Grão-Pará North, Centre and South. Some haphazard collections were made in Camp Curuá (Estanífera) of the mining company rio Tinto. The area is drained, from W to E, by the headwaters of the Mapuera and Trombetas rivers, and by the Paru de Oeste, Cuminapanema, Curuá and Maicuru rivers.

Camp ESEC Grão-Pará North (N 1° 17' 7.51" W 58° 41' 45.24") was situated in the NW part of the unit, in the municipality of Oriximiná, close to the border with Guyana in the Acarai Mountains (Figure 11), at an altitude of 500 m on a hill. The area is covered by terra-firme forest ('Floresta Ombrófila Densa Submontana') and is very hilly, with steep slopes leading down to creeks with clear, transparent water. Relatively small rock outcrops occur sparsely in the area, completely covered by forest. From the camp, trail 1 ran SE for 4.35 km, first descending into a valley at 400 m, then climbing out of the valley up a spur of a hill to 600 m, down again to a second valley at about 460 m and up another hillside to 500 m. Trail 2 ran S for 900 m and then SW, following a spur of the hill on which the camp was situated, at the end slightly going down, and to the west, after 2.8 km joining trail 3 at km 2.5. Trail 3 ran SSW for 4.



km, descending into a valley at 400 m altitude, following a creek, at km 2.5 it was joined by trail 2 from the east. Trail 4 started at km 0.9 of trail 2 where it split off to SSE for 2.1 km, down from the camp into a valley at 400 m altitude. Pitfalls were placed in trail 1 at 150 m, 250 m, 450 m, 650 m, 1,800 m, 2,000 m, 2,150 m and 2,260 m from camp; and in trail 3 at 300 m, 550 m, 750 m, 950 m, 1,100 m, 1,250 m, 1,450 m, and 1,650 m from camp. In trails 2 and 4 no pitfalls were placed. Altitudes varied between 350 and 600 m above sea level. ESEC Grão-Pará North was the aim of the fourth expedition, between August 25 and September 11, 2008, during the dry season.

Camp ESEC Grão-Pará Centre (N 0° 37' 49.01" W 55° 43' 42.60") was situated in the municipality of Óbidos, in the northern part of the eastern half of the ESEC, close to the southern border of the TI Tumucumaque. It was at an altitude of 400 m in a transition zone at the W margin of a large island of terra-firme forest ('Floresta Ombrófila Densa Submontana') within a large savanna enclave. The savanna area consisted of hilly terrain (300 to 500 m), with many areas of rock outcrops (from horizontally flat to curved and steep) and with a vegetation of shrubs and low forest (Figure 12), sometimes interrupted by grassy areas with isolated trees (e.g. *Curatella*) (Figure 13). Creeks in the savanna were rare. One encountered was a deep (3 m) gully with vertical banks and did not contain any water. Another creek arose at the base of a large complex of rock outcrops and contained clear, transparent water. This creek ran at the border between open rock outcrop and savanna forest and formed some deeper pools connected by shallower areas. Open rock outcrops were generally wet and retained water in crevices and under loose rocks, well after rains had stopped (Figure 14). Hillsides were generally steep. The forest was terra-firme forest with large trees and a high canopy at about 30 m. The forest island was traversed by a large creek with (at the time shallow) clear, transparent water, in some places forming deeper pools. The transitional forest between forest and savanna consisted of small, slender-stemmed, low trees with some

larger trees interspersed; this vegetation was characterized by the botanists as 'cerradão'. From camp, trail 1 ran N for 5 km through terra-firme forest and slightly undulating terrain, crossing the creek in the forest island at about 1 km from camp. Trail 2 ran NE for 5 km, reaching the E edge of the forest, first dropping into the bed of the aforementioned creek, than steeply climbing up to a plateau at 500 m and than dropping again to a level of 400 m. Trail 3 ran S for 4.4 km, generally through savanna over steep hills, but at 700 m and at 2.5 km entering narrow areas of forest with creeks. Trail 4 ran roughly W for 5 km through open savanna, savanna forest and open rock outcrops, twice crossing narrow areas of terra-firme forest without creeks. The first part of the trail was in an area with steep hills, and after 1450 m descended into a relatively flat area. Pitfalls in trails 1 and 2 were placed at 500 m, 750 m, 1,000 m and 1,250 m from camp, all in forest. In trail 3 (savanna) they were placed at 200 m, 450 m, 750 m and 900 m. In trail 4 (savanna) they were placed at 400 m, 800 m, 1,240 m and 1,440 m. Altitudes varied between 310 and 450 m above sea level. ESEC Grão-Pará Centre was inventoried between January 10 and 31, 2009, during the early rainy season.

Camp ESEC Grão-Pará South (S 0° 9' 55.76" W 55° 11' 11.04") was situated in the municipality of Alenquer, in the SE part of the ESEC, close to its border with FLOTA Paru, and only 6 km NW of the Base Curuá (Estanífera) of Rio Tinto, at an altitude of 300 m. It was situated in a wide creek valley surrounded by hills. The area, according to vegetation maps, was covered by 'Floresta Ombrófila Densa Submontana', but along creeks, around a lake present W of the camp, and on top of a plateau, the forest was clearly different from the terra-firme forest covering the hills. In the creek valleys and near a lake were patches of 'açaiçal', açai forest dominated by *Euterpe* palms in shallow water. The forest along creeks in low-lying areas generally was open, with a muddy surface and pools, tufts of grasses, few large trees and many thin trees; it was considered as 'igapó forest' by the botanists. From

the camp, trail 1 ran NNE for 5 km, first crossing the side of a hill, than dipping into a partly inundated creek valley, before starting a long climb up to a plateau that was covered by a low type of forest (canopy at 10 m) that consisted of closely growing small trees with thin stems (called 'cerradão' by the botanists). Only few larger trees were present. Trail 2 ran E for 5 km, first through a level creek valley, but after 2 km climbing a ridge and going down again to the next creek valley. Trail 3 ran SE for 5 km, through a swampy area along a creek, after 3 km continuing on the lower part of a hill, more or less parallel to a creek valley. Trail 4 ran W for 3.4 km through a creek valley, in the first 50 m crossing a very dense liana forest, than entering igapó forest and crossing a large deep creek, with steep sides and water with much organic particles, several times. Along this trail there were several ponds (1.5 m deep) in the forest. At the end the trail reached a sizeable open lake and ran around it. The lake on most sides was bordered by palms (*Attaleia spinosus*, *A. maripa*, *Mautitia flexuosa*), but on its north side was bordered by a sloping area with terra-firme forest that reached the water margin. The centre of the lake was covered by a vegetation of Cyperaceae and grasses, along its edges there were waterlilies. The water was very clear and transparent (at least 2 m view). A creek flowed out of the lake on the east and passed the camp about 3 km downstream. On the west a small creek, completely overgrown with vegetation, flowed into the lake. The altitude varied from 300 m in the creek valleys to 450 m on top of the plateaus. Pitfalls were only placed in trails 1 (10 sets) and 3 (6 sets). In trail one they were at 1,550 m, 1,650 m, 1,800 m, 2,000 m, 4,000 m, 4,100 m, 4,200 m, 4,300 m, 4,400 m and 4,500 m. The first four were in terra-firme forest between 320 and 380 m altitude, the last six, from 4,000 m on, were in cerradão forest at 420 m altitude. Those in trail 3 were at distances of 800 m, 1,230 m, 1,280 m, 1,500 m, 1,650 m, and 1,750 m from camp. No pitfalls were placed in trails 2 and 4 because large areas of those trails were inundated creek valley. The third expedition

targeted ESEC Grão-Pará South between June 6 and 21, 2008, towards the end of the wet season.

Base Curuá (Estanífera) of mining company Rio Tinto (S 0° 13' 16.5" W 55° 09' 45.0"), in the municipality of Alenquer, although not an area that was systematically collected, like the ones described before, is shortly mentioned because some species were collected here that were not collected elsewhere. It is situated 8 km SE of ESEC Grão-Pará South, on top of a plateau (450 m), with an airstrip and an area where semi-permanent barracks have been mounted for personnel of Rio Tinto working in the area. The area around the campsite and near the airstrip embarking site consists of a low forest (canopy 5 m) with many narrowly spaced thin-stemmed trees (Figure 15). This forest was characterized by the botanists as 'cerradão'. Along a trail to the rio Curuá there was a small open rock savanna, with large areas of bauxite rock on the surface and a sparse vegetation of herbs, Ananas and shrubs.

## LOGISTICS

All expeditions started in Belém and went to Santarém by commercial flight, except the expedition to ESEC Grão-Pará North which flew from Belém to Boa Vista, Roraima. The expedition to FLOTA Faro used a boat as base camp. The expedition to ESEC Grão-Pará North traveled by bus from Boa Vista to Caroebe (still in Roraima) and from there by helicopter to base camp ESEC Grão-Pará North. In the other five expeditions participants were transported from Santarém by one-engine planes to either Camp Curuá (Estanífera) of Rio Tinto, to airstrip '13 de maio' (REBIO Maicuru) or to Monte Dourado (FLOTA Paru). From those places transport to the camps was by helicopter, except for the expedition to FLOTA Paru, which from Monte Dourado went by bus to the rio Paru and crossed the river by canoes with an outboard motor. Base camps, trails and pitfalls were prepared shortly before arrival of the scientific participants. Camp size tended to increase during the year that expeditions were held, from 210 m<sup>2</sup> in FLOTA Trombetas, to 350 m<sup>2</sup> during the last expedition (ESEC Grão-Pará Centre). Helicopter time amounted to 287:10 h, small airplanes time to 334:53 h.



## COLLECTING AND PREPARATION OF HERPETOFAUNA

Reptiles and amphibians were collected by means of two complementary methods, *viz.* active sampling (AS), which combined two techniques: time constrained searches (TCS) and visual and audio encounter surveys (VES) (no recordings of calls were made); and passive sampling by pitfall traps with drift fences (PD). Systematic collecting took place in all seven expedition areas described above. Camp Curuá (Estanífera) is not a major collecting site, no systematic sampling was done there and it has not been included in any calculations, except for total number of species in the whole area, since a few species have been collected only there. For the same reason it appears in the table presented in the Appendix 1.

All trails and most pitfalls were georeferenced, but we here only present the coordinates of the centrally located campsites (mark 0 of the trails of every expedition).

Active sampling (Crump & Scott, 1994; Scott, 1994; Ribeiro-Junior *et al.*, 2008) consisted in actively searching for animals during day and night, along marked trails and in different habitats, noting time spent (unit of collecting = person.hour). AS is important for a general inventory, considering both taxonomic coverage (pitfalls are only adequate for some groups) and coverage of different habitats. This method requires trained personal, and, even with experienced collectors, is subject to personal bias, which causes problems to compare areas or studies done by this method. Another unequal factor, in the present case, was that three expeditions had two herpetologists and four had three, and thus collecting effort in active sampling was not uniform for all expeditions.

Pitfall traps with drift fences (Corn, 1994; Cechin & Martins, 2000; Ribeiro-Junior *et al.*, 2008): each trapping unit consisted of four buckets (pitfalls) of 60 l each, that were dug into the ground with their rim flush with the surface, positioned in a Y, with the central bucket connected to the three peripheral ones by eight meters of black plastic sheet with a height of 60 cm. A total of 16 trapping units was

used, positioned in two, three or four groups (each group in a different trail), depending on the local conditions (type of substrate, rocks, flooding, logistics). Within each group, trapping units generally were positioned at distances of 250 m from each other, but in FLOTA Trombetas they were partly 125 m apart, and in some cases greater distances were used in order to also sample different vegetations or because of physical problems encountered in the terrain. Pitfalls were checked once a day during the entire sampling period. Trapping units were installed in the week before the start of each expedition and removed at the end of each expedition. Collecting with pitfalls has the advantage to be independent of collector and they collect species (generally (semi) fossorial) that are only rarely caught during active collecting. On the other hand, pitfalls are directed to leaf litter and terrestrial species, although some arboreal species that come to the ground are also collected with certain regularity; other groups, especially arboreal/climbing and aquatic species, large terrestrial species and medium-sized to large snakes, and a number of amphibians that are able to climb or jump out of the buckets, are rarely collected. Besides, it is not always possible to use pitfalls; flooded or rocky areas, or areas far from basecamp, can not be sampled by this method, because of the impossibility to install the buckets in the first areas, and the impossibility to check the pitfalls every day in the last case.

Specimens collected by the fieldworkers or by other researchers were considered as occasional encounters.

For each specimen observed by the herpetologists and/or collected the following data (if applicable) were annotated: identification, locality (GPS or distance in trail), habitat, microhabitat, day, time, and name of collector/observer. Collected animals' standard measurements and weight were taken, and when possible the sex was determined. In many cases notes of life colouration were made, and digital photographs and tissue samples (liver or muscle) for molecular studies were taken.

Collected specimens were euthanized by an overdose of veterinary anesthetic, fixed in formaldehyde



4% (one part of commercial formaldehyde 37% and nine parts of water) for maximally 24 hours, labeled with field numbers (a field series 'CN' [Calha Norte] was created for all specimens collected during the expeditions), and preserved in alcohol 70%, except tadpoles that were preserved in 4% formaldehyde (Franco *et al.*, 2002). Tissue samples were preserved in absolute alcohol, maintained at environmental temperature in the field and transferred to a freezer after arrival in the museum. Even though we refer here to number of specimens per site, it is important to keep in mind that each site covers a few square kilometers and different environments. Thus specimens dealt with as coming from one site may have been collected up to 8,000 m apart. All material was deposited in the herpetological collection of Museu Paraense Emílio Goeldi (MPEG) in Belém, Pará, Brazil (Appendix 2). Collections were made under licence 001/2008 of the Secretaria Estadual do Meio Ambiente (SEMA-PA).

Collecting effort, number of specimens collected per group, and rarefaction curves for amphibians and reptiles separately, are presented for each major collecting site (thus, not for Estanifera). Species rarefaction curves were constructed with the help of the program EstimateS version 7.5 (Colwell, 2005), on the basis of grouped AS and PD collections, considering as sampling unit each collecting day – they represent the cumulative number of species against the increase of collecting effort, obtained after 50 randomizations. Species composition in the seven sites is compared, using the Biogeographic Similarity Coefficient (Duellman, 1990), equivalent to  $2C/(N_1 + N_2)$ , where  $N_1$  = number of species in locality 1,  $N_2$  = number of species in locality 2, and  $C$  = number of species common to both localities. We present a gross estimation of relative abundance of amphibians and reptiles, for each site and for each group as a whole, based on numbers of specimens caught by pitfall traps and all those collected by active searches. Species observed but not collected are added to the graphs as presenting '0' (zero) specimens.

Species were considered endemic for the Guianan Region mainly based on Hoogmoed (1979b) (reptiles and amphibians), Señaris & MacCulloch (2005) and Duellman (1999) for amphibians, and Avila-Pires (2005) for reptiles.

For familial and generic nomenclature we adhere to the nomenclature used before the Faivovitch *et al.* (2005), Frost *et al.* (2006), and Grant *et al.* (2006) papers, because we are not convinced that the wholesale changes in nomenclature proposed by these authors are really necessary and correct. We prefer to await further independent studies that corroborate the alterations those publications proposed. We have made an exception for the former species of *Colostethus* in the Guianan Region with a mid-lingual papilla, which are clearly recognizable morphologically and now are named *Anomaloglossus* (Grant *et al.*, 2006). Also, we have not yet taken into account the changes proposed for the genus *Eleutherodactylus* by Heinicke *et al.* (2007) and by Hedges *et al.* (2008). Discussion about these issues is still going on, and by maintaining the pre-2005 names we keep the relation with older (= pre-2005) literature, thus facilitating the work of conservationists and managers of natural areas.

Two recent publications, Zaher *et al.* (2009) and Vidal *et al.* (2010), proposed changes in the classification of South American Colubrid snakes. Because of the short time to properly evaluate them (when they appeared this paper already had been completed), we preferred not to incorporate their changes here.

We tried to identify material of the *Bufo granulosus* complex with Narvaes & Rodrigues (2009), but had serious problems trying to separate the species recognized by these authors in the area north of the Amazon River (Gorzula & Señaris (1999) who also doubted the validity of different taxa in Venezuelan Guiana). We doubt whether all taxa recognized by Narvaes & Rodrigues (2009) are real entities and for the time being we have adhered to the use of *Bufo granulosus* for the medium-sized granular toad with dorsally directed nostrils, occurring north of the Amazon, realising that e.g. *Bufo mirandaribeiroi* Gallardo, 1965 is a good taxon.



## COLOUR PHOTOS

We present colour photos of some of the species found during the expeditions (Figures 16-75). They generally appear in the same sequence as used in Appendix 1.

## RESULTS AND DISCUSSION

### COLLECTING EFFORT AND SPECIES RICHNESS

Collecting effort per expedition for each method is shown in Table 1, which also refers to the main habitats found in each site. Taking all areas together we registered 80 species of amphibians (77 frogs and three caecilians) and 95 species of reptiles (36 species of lizards, three of amphisbaenians, 49 species of snakes, five species of chelonians and two species of caiman). The large lizard *Tupinambis teguixin* (Linnaeus, 1758) (jacuraru or teiu) seems to have been seen in some areas, but not by herpetologists. Because of doubts remaining about the identification, we have not included this species in our list, although it is expected to occur throughout the area (see below). Comparing these numbers with what is expected to occur in the area based on the literature (see introduction), we see that lizards are well represented, followed by frogs, while caecilians, amphisbaenians, snakes and chelonians are well below the expected richness, being clearly underrepresented in the samples. Chelonians, living mainly in aquatic environments, need special collecting techniques, not used in the surveys. Caecilians, partially aquatic and partially fossorial, and amphisbaenians, which are mainly fossorial, are only sporadically found, even when digging extensively for them. Snakes have usually secretive habits and are known to need long periods for inventorying them satisfactorily in tropical rainforests.

Table 2 shows the number of species per expedition. The highest number of amphibians (36 species) was registered for ESEC Grão-Pará South and the lowest number (21 species) for the FLOTA Faro. The highest number of reptiles (42 species) was collected in ESEC Grão-Pará North, and the lowest number (27 species) in

ESEC Grão-Pará South. However, none of the rarefaction curves, calculated for each site, separately for amphibians and reptiles, reached the asymptote (Figure 76), indicating that more species are expected to occur in each area. The need of prolonged effort to adequately survey the herpetofauna is well documented in the literature (e.g., Duellman, 1978; Myers & Rand, 1969). Duellman (2005) reported that, in Cusco Amazónico, it took 442 person-days to record 89% of the species of anurans, 81% of the lizard species, and 79% of the snake species present in the area. Ribeiro-Junior *et al.* (2008) also showed the necessity of a large collecting effort, using multiple techniques, for an adequate representation of the herpetofauna. Results of collections vary due to several factors, among others time of the year, meteorological circumstances during the expedition, micro- and meso-habitats sampled, and in the case of active collecting the collectors and the number of collectors. For example, in ESEC Grão-Pará South, the larger number of amphibian species was partly due to the presence of ponds and a lake that provided good conditions for amphibians, especially Hylidae. Although similar habitats have not been encountered in the other sampled areas, it is very unlikely that they do not occur in all conservation units. Besides, it should be remembered that some amphibians use for reproduction temporary pools that only form during the rainy season, and some species have explosive reproduction which only lasts a few days. All these factors influence the number of species found during the limited period of an expedition.

Comparing species richness obtained for each area with number of specimens collected (Figure 77), the studied sites in FLOTA Trombetas, REBIO Maicuru and ESEC Grão-Pará show an increasing number of species with increasing number of specimens collected. Richness in FLOTA Faro and FLOTA Paru, on the other hand, is proportionally less in relation to the number of specimens collected. These two areas are situated along large rivers bordered by flooded forest, a habitat that usually has less species than terra-firme forest (also present in these sites). However, the savanna vegetation

Table 1. Collecting effort per method, and main habitats at each studied site. AS = Active Search, PD = pitfalls with driftence.

	FLOTA Faro	FLOTA Trombetas	ESEC Grão-Pará North	ESEC Grão-Pará Centre	ESEC Grão-Pará South	REBIO Maicuru	FLOTA Paru
AS daytime (person.hours)	72	100	180	172 (74f, 98c)	134	162	78
AS night (person.hours)	72	18	20	88 (40f, 48c)	36	58	19
PD (trap.nights)	192	237	240	240 (120f, 120c)	208	240	208
Habitats	Terra-firme forest (3 trails); flooded forest (1 trail); disturbed areas	Terra-firme forest with creeks (some areas with many <i>Cecropia</i> ); an open area with rock outcrop	Terra-firme forest, with steep slopes and creeks	Terra-firme forest (f) and cerrado (c) with rock outcrops	Terra-firme forest with creeks bordered by flooded areas; an extensive lake bordered by palms	Terra-firme forest and (dry) flooded forest; small beach along margin of river	Terra-firme forest with creeks and different degrees of perturbation (3 trails); (dry) flooded forest along margin of river (1 trail)

Table 2. Number of species per expedition and per taxonomic group. The column "Grão-Pará (Estanífera)" refers to the Rio Tinto basecamp (rio Curuá), where only occasional collections were made, but where two species were collected that were found nowhere else in ESEC Grão-Pará.

	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanífera)	REBIO Maicuru	FLOTA Paru	Total
AMPHIBIA	21	30	24	32	36	1	31	24	<b>80</b>
REPTILIA	35	32	42	29	26	1	34	29	<b>95</b>
lizards and amphisbaenians	20	16	24	15	14	0	21	18	<b>39</b>
snakes	12	13	16	12	10	1	10	10	<b>49</b>
chelonians	2	2	1	1	2	0	2	1	<b>5</b>
cayman	1	1	1	1	0	0	1	0	<b>2</b>
<b>TOTAL</b>	<b>56</b>	<b>62</b>	<b>66</b>	<b>61</b>	<b>62</b>	<b>2</b>	<b>65</b>	<b>53</b>	<b>175</b>

that covered part of ESEC Grão-Pará Centre also harbors a lower number of species than terra-firme forest, but in spite of that, relative richness was proportional to that in other sites. The two FLOTA, Faro and Paru, also had in common areas of disturbed and secondary forests, and one possibility is that the lower relative richness of species in these two areas is a result of environmental disturbance. Habitat alteration may have caused the disappearance, or a population reduction (making them more difficult to be captured), of a number of species.

## SPECIES COMPOSITION

A complete list of species (including authors and years), expeditions during which they were collected and basic biological data are presented in the Appendix 1. Looking at the herpetofauna as a whole, only 5.7% of the species were found in all seven sites, while 43.4% were found in only one of the sites (Table 3). Lizards and amphisbaenians, as a group, showed the most even distribution in the samples, with almost 18% of the species captured in all






Figure 1. Map of the eastern part of the Guianan Region, showing Indian Territories and protected areas in Brazil (Amapá, northern Pará, eastern Amazonas and Roraima), French Guiana, Suriname and eastern Guyana.






Figure 2. Map of northern Pará with localities inventoried (red dots), and areas where inventories were made before the Calha Norte Project (light blue dots). The purple area around point 2 is savanna. 1. ESEC Grão-Pará North, 2. ESEC Grão-Pará Centre, 3. REBIO Maicuru, 4. ESEC Grão-Pará South and Estanifera, 5. FLOTA Trombetas, 6. FLOTA Paru, 7. FLOTA Faro, 8. FLONA Saracá-Taquera, 9. Rio Nhamundá, Sítio Céu Estrelado-Rio Trombetas, Cruz Alta, 10. Monte Alegre, 11. Jari.




Figure 3. Mean annual precipitation for northern Pará.




Figure 4. Mean annual temperatures for northern Pará.



Figure 5. FLOTA Faro, view of inundated igapó forest. Photo: Luciano F.A. Montag.



Figure 6. FLOTA Trombetas, trail 3, terra-firme forest with large rocks. Photo: MSH.



Figure 7. FLOTA Trombetas, heliporto with open lajedos and dry-adapted vegetation (cacti, bromeliads). Photo: MSH.



Figure 8. REBIO Maicuru, terra-firme forest with flowering trees (*Tabebuia* sp.) in the dry season. Photo: MSH.



Figure 9. REBIO Maicuru, rio Ipitinga. Photo: MSH.



Figure 10. FLOTA Paru, looking south along the rio Paru. Photo: MSH.





Figure 11. ESEC Grão-Pará North, general outlook over the Acarai Mountains. Photo: TCSAP.



Figure 12. ESEC Grão-Pará Centre, trail 3, km 0.9, open rock outcrop and savanna vegetation. Habitat of *Gymnophthalmus* cf. *underwoodi*. Photo: MSH.



Figure 13. ESEC Grão-Pará Centre, pitfall 9, installed in savanna area of trail 3. Photo: MSH.



Figure 14. ESEC Grão-Pará Centre, trail 4, km 0.9, open, wet rock outcrops surrounded by savanna vegetation. Habitat *Leptodactylus myersi*. Photo: MSH.



Figure 15. Camp rio Curuá (Estanifera) of Rio Tinto mining company. The 'cerradão' vegetation where *Rana palmipes* and *Physalaemus ephippifer* were found in small poodles. Photo: MSH.



Figure 16. *Cochranella* sp. (CN2363), ESEC Grão-Pará Centre. Photo: MSH.



Figure 17. *Atelopus hoogmoedi*, male (CN2086), ESEC Grão-Pará Centre. Observe difference in pattern with male. Difference is not sex-related. Photo MSH.



Figure 18. *Atelopus hoogmoedi*, female (CN2102), ESEC Grão-Pará Centre. Photo: MSH.



Figure 19. *Bufo margaritifer*, female (CN421), FLOTA Trombetas. Photo: MSH.



Figure 20. *Bufo margaritifer*, male (CN426), FLOTA Trombetas. Photo: MSH.



Figure 21. *Epipedobates cf. guayanensis* (CN394), FLOTA Trombetas. Photo: MSH.



Figure 22. *Epipedobates hahneli* (CN495), FLOTA Trombetas. Photo: MSH.



Figure 23. *Dendrobates tinctorius*, blue variety (September 12, 2008, not collected) ESEC Grão-Pará North, courtship behaviour. Photo: TCSAP.



Figure 24. *Hyla dentei* (CN 1492), REBIO Maicuru. Photo: MSH.



Figure 25. *Hyla gaucheri* (CN825), ESEC Grão-Pará South. Photo: MSH.



Figure 26. *Hyla punctata* (CN1994), FLOTA Paru. Photo: MSH.



Figure 27. *Phyllomedusa tomopterna* (CN2134), ESEC Grão-Pará Centre. Photo: TCSAP.



Figure 28. *Scinax garbei* (CN1629), REBIO Maicuru. Photo: TCSAP.



Figure 29. *Scinax proboscideus* (CN712), ESEC Grão-Pará South. Photo: MSH.



Figure 30. *Rana palmipes* (CN600-601), Estanifera (Acampamento Curuá). Photo: MSH.



Figure 31. *Adenomera andreae* (CN), REBIO Maicuru. Photo: MSH.



Figure 32. *Adenomera hylaedactyla* (CN2113), ESEC Grão-Pará Centre. Photo: MSH.



Figure 33. *Ceratophrys cornuta* (CN 799), ESEC Grão-Pará South. Note difference in colour with other specimen depicted. Photo: MSH.



Figure 34. *Ceratophrys cornuta* (CN 805), ESEC Grão-Pará South. Note difference in colour with other specimen depicted. Photo: MSH.




Figure 35. *Leptodactylus knudseni* (CN2195), ESEC Grão-Pará Centre. Photo: MSH.



Figure 36. *Leptodactylus longirostris* (CN2110), ESEC Grão-Pará Centre. Photo: MSH.



Figure 37. *Leptodactylus myersi* (April 26, 2008, not collected), FLOTA Trombetas. Photo: MSH.



Figure 38. *Leptodactylus pentadactylus* (CN920), ESEC Grão-Pará South. Photo: MSH.



Figure 39. *Lithodytes lineatus* (CN453), Flota Trombetas. Photo: MSH.



Figure 40. *Physalaemus ephippifer* (CN562), FLOTA Trombetas. Photo: MSH.



Figure 41. *Hamptophryne bolivianus* (CN796), ESEC Grão-Pará South. Photo: MSH.



Figure 42. *Otophryne pyburni* (CN1358), ESEC Grão-Pará North. Photo: TCSAP.



Figure 43. *Synapturanus mirandaribeiroi* (CN373), FLOTA Trombetas. Photo: MSH.



Figure 44. *Pipa snethlageae* (CN319), FLOTA Faro. Photo: W.A. Rocha.



Figure 45. *Caecilia tentaculata* (CN 2138), ESEC Grão-Pará Centre. Photo: MSH.



Figure 46. *Rhinatrema* sp. n. (CN1088), ESEC Grão-Pará North. Photo: TCSAP.



Figure 47. *Coleodactylus amazonicus* (CN530), FLOTA Trombetas. Photo: MSH.



Figure 48. *Lepidoblepharis heyerorum*, male (CN248), FLOTA Faro. Photo: W.A. Rocha.



Figure 49. *Amapasaurus tetradactylus* (CN 362), FLOTA Trombetas. Photo: MSH.



Figure 50. *Bachia flavescens* (CN791), Grão-Pará South. Photo: MSH.



Figure 51. *Cercosaura ocellata*, male (CN2028), FLOTA Paru. Photo: MSH.



Figure 52. *Gymnophthalmus* cf. *underwoodi* (CN2225) ESEC Graõ-Pará Centre. Photo: MSH.





Figure 53. *Iphisa elegans* (CN1100), ESEC Grão-Pará North. Photo: TCSAP.



Figure 54. *Leposoma guianense*, male (CN1095), ESEC Grão-Pará North. Photo: TCSAP.



Figure 55. *Neusticurus rudis* (CN168), ESEC Grão-Pará North. Photo: TCSAP.



Figure 56. *Tretioscincus agilis* (CN1232), ESEC Grão-Pará North. Photo: TCSAP.



Figure 57. *Kentropyx striata* (CN1247), ESEC Grão-Pará Centre. Photo: MSH.



Figure 58. *Mabuya nigropunctata* (January 18, 2009, not collected), ESEC Grão-Pará, trail 1 km 0.65. Photo: MSH.





Figure 59. *Anolis auratus* (CN2973), ESEC Grão-Pará Centre. Photo: MSH.



Figure 60. *Anolis chrysolepis*, male (CN295), FLOTA Faro. Photo: W.A. Rocha.



Figure 62. *Amphisbaena vanzolinii* (CN639), ESEC Grão-Pará South. Photo: MSH.



Figure 61. *Uranoscodon superciliosus* (CN2300), ESEC Grão-Pará Centre. Photo: MSH.



Figure 63. *Mesobaena* sp.nov. (MPEG 24854, CN 7), FLOTA Faro. The animal was cut while digging during installing pitfalls. A small middle section could not be recovered. The colours are those just after death of the animal. Photo: MSH.



Figure 64. *Leptotyphlops albifrons* (CN2069), ESEC Grão-Pará Centre. Photo: Pedro L.V. Peloso.



Figure 65. *Corallus caninus* (CN973), ESEC Grão-Pará North. Photo: TCSAP.



Figure 66. *Apostolepis nigrolineatus* (CN 1317), ESEC Grão-Pará North. Photo: TCSAP.



Figure 67. *Atractus badius* (CN 804). ESEC Grão-Pará South. Photo: MSH.



Figure 68. *Chironius scurrulus* (CN1930), FLOTA Paru. Photo: MSH.



Figure 69. *Pseudoboa newwiedii* (CN 2218), ESEC Grão-Pará Centre. Photo: MSH.



Figure 70. *Thalesius viridis* (CN 1292). ESEC Grão-Pará North. Photo: TCSAP.



Figure 71. *Micrurus averyi* (CN 1086). ESEC Grão-Pará North. Photo: TCSAP.



Figure 72. *Micrurus hemprichii* (CN2287), ESEC Grão-Pará Centre. Photo: MSH.



Figure 73. *Micrurus paraensis* (CN793), ESEC Grão-Pará South. Photo: MSH.



Figure 74. *Chelonoidis carbonaria*, juv. (CN622), ESEC Grão-Pará South. Photo: MSH.



Figure 75. *Chelonoidis denticulata*, male (June 13, 2008, not collected), ESEC Grão-Pará South trail 2, km 0.7. Photo: MSH.




Figure 76. Rarefaction curves for each studied site calculated based on cumulative number of species in relation to the number of collecting days, showing the 95% confidence interval lines. Gray = reptiles, black = amphibians, X axis = number of samples (collecting days), Y axis = cumulative number of species.




Figure 77. Relation between number of species registered during each expedition and number of specimens collected. Numbers above the squares refer to the studied sites as they appear in Fig. 2. 1 = ESEC Grão-Pará North, 2 = ESEC Grão-Pará Centre, 3 = REBIO Maicuru, 4 = ESEC Grão-Pará South, 5 = FLOTA Trombetas, 6 = FLOTA Paru, 7 = FLOTA Faro.

Table 3. Distribution of species (number and percentage) in relation to the number of studied sites in which they were found. Chelonians and caimans are included in the total numbers but are not shown separately.

Number of sites	Amphibians		Lizards and amphisbaenians		Snakes		Total	
	Number of species	%	Number of species	%	Number of species	%	Number of species	%
7	3	3.8	7	17.9	0	0	10	5.7
6	4	5.1	3	7.7	0	0	7	4.0
5	7	8.9	1	2.6	0	0	9	5.1
4	6	7.6	2	5.1	2	4.1	10	5.7
3	10	12.7	6	15.4	7	14.3	25	14.3
2	13	16.5	9	23.1	14	28.6	37	21.1
1	36	45.6	11	28.2	26	53.1	76	43.4

sites, and only 28% in only one site. Of amphibians, about 46% were found in only one site, and of snakes 53%. The maximum number of sites a species of snake occurred in was four, once more showing the haphazardness of finding these animals. The low number of amphibians common to all sites is partially linked to differences in habitats available

and surveyed, as well as in period of the year. But certainly for all groups part of the differences is due to chance and should decrease as collecting effort increases. The same applies when we look at the Biogeographic Similarity Coefficient (BSC) (Table 4). This coefficient, that represents the proportion of species common to two areas, in relation

Table 4. Total number of species per site (in bold, diagonal line), number of species in common (above diagonal line), and Coefficient of Biological Similarity (below diagonal line) between each pair of sites.

	FLOTA Faro	FLOTA Trombetas	ESEC Grão-Pará North	ESEC Grão-Pará Centre	ESEC Grão-Pará South	REBIO Maicuru	FLOTA Paru
FLOTA Faro	<b>56</b>	28	26	22	24	26	22
FLOTA Trombetas	0.47	<b>62</b>	28	30	31	30	31
ESEC Grão-Pará North	0.43	0.44	<b>66</b>	23	25	29	27
ESEC Grão-Pará Centre	0.38	0.49	0.36	<b>61</b>	28	25	27
ESEC Grão-Pará South	0.41	0.50	0.39	0.46	<b>62</b>	29	29
REBIO Maicuru	0.43	0.47	0.44	0.40	0.46	<b>65</b>	34
FLOTA Paru	0.49	0.54	0.45	0.47	0.50	0.58	<b>53</b>

to the total number of species in both areas, is lowest (0.36) between ESEC Grão-Pará Centre and ESEC Grão-Pará North. While the northern part of this ESEC lies in the Acarai Mountains, covered by rainforest in an extremely hilly area, the site in the central part of this conservation unit presents a large isolated area of open vegetation (savanna) (with its specific fauna), with a patch of more or less isolated forest in its interior, which explains part of the difference found between these two areas. In addition, the northern sector was sampled in August-September, thus during the dry season, while sampling in the central sector occurred in January, in the early rainy season, which probably also accounts for part of the differences found. Going to the other extreme, FLOTA Paru and REBIO Maicuru were the most similar areas (BSC 0.58). These two areas were sampled within a period of three months and both have seasonally flooded areas (igapó) influenced by a river, which may explain their larger similarity. But again, the expectation is that these coefficients, between all these areas, will become larger (indicating more similar herpetofaunas) as a

better representation of the herpetofauna, from all around the year, is obtained.

Some species however are restricted to special habitats, which may account for real differences between sites. Thus, for example, *Hyla wavrini*, a várzea and igapó species, was only collected in FLOTA Faro, whereas in other localities *Hyla boans*, a very similar species that occurs along creeks and rivers in terra-firme rainforest or in gallery forest (Hoogmoed, 1990a), was collected. It is possible that these two species are mutually exclusive. *Atelopus hoogmoedi* and the two species of Centrolenidae collected depend on the presence of (relatively) clear, running water with rapids in terra-firme forest, and only were recorded for ESEC Grão-Pará Centre (with unidentified Centrolenid tadpoles collected in ESEC Grão-Pará North and South). Although they are without doubt present in other areas of CNP, their distribution is limited to specific habitats. The same is true for *Leptodactylus myersi*, restricted to large, open rocky slabs, either surrounded by terra-firme forest (FLOTA Trombetas) or in savanna areas (ESEC

Grão-Pará Centre). Savanna enclaves, as encountered in ESEC Grão-Pará Centre, have a herpetofauna that is largely different from that of forested areas, among which *Adenomera hylaedactyla*, *Leptodactylus longirostris*, *Anolis auratus*, *Kentropyx striatus*, *Gymnophthalmus cf. underwoodi* and *Pseudoboa newwiedii*. These species are restricted to these enclaves and, some of them, to river beaches along the Amazon River.

The caimans *Paleosuchus trigonatus* and *P. palpebrosus* (Cuvier, 1807) are small and live in creeks in the forest, while *Caiman crocodilus* occurs in rivers and larger creeks which are not completely roofed over by forest canopy. *Paleosuchus trigonatus* was found in creeks in FLOTA Trombetas and ESEC Grão-Pará North and Centre. *Paleosuchus palpebrosus* was not encountered during the expeditions, but it is known from Oriximiná, Trombetas River (Medem, 1983), and is probably present in other areas of CNP as well. *Caiman crocodilus* was found in FLOTA Faro and in REBIO Maicuru, and it is possible that it also occurs in the Jari, Paru and Trombetas rivers, but not far from the main course of these larger rivers.

*Dendrobates tinctorius* (blue variant) only was encountered in the Acarai Mountains near the border with Guyana, in the northern part of ESEC Grão-Pará, where it was quite numerous. However, there are also records of this species from Porto Trombetas (blue variant) (material in MPEG) and in Monte Dourado (variant with large dorsal yellow patch and lines) (material in MPEG), where they occur in some forest localities, but not in others. In this case, the distribution of the species seems to be patchy, but it is not clear which environmental parameters are important to decide in which parts of an area it occurs.

The southeastern portion of ESEC Grão-Pará, including the studied points in the centre and south of the reserve, presents patches of forest consisting of small, slender-stemmed, low trees with some larger trees interspersed, known as 'cerradão'. No amphibian or reptile species was found only in this type of vegetation, where at least some of the forest species are present.

## COMPARISON WITH A SITE IN SOUTHERN GUYANA

As pointed out above, publications dealing with the fauna of northern Pará are scarce. However, there is one paper (Señaris *et al.*, 2008) that deals with a site in southern Guyana that is only about 50 airline kilometers southwest of our collecting site ESEC Grão-Pará North. This is much nearer than ESEC Grão-Pará North is to any of the other six studied localities in CNP. It could be expected that the herpetofaunas of these two areas would be very similar. Although the methodology of Señaris *et al.* (2008) differs considerably from ours (they did not use pitfalls and driftfences, only opportunistic surveys and Visual Encounter Survey) it seems worthwhile to make some comparisons. Señaris *et al.* (2008) reported 26 species of amphibians (25 frogs, one *Gymnophiona*) and 34 species of reptiles (12 lizards, one amphisbaenian, 16 snakes, three chelonians, two caiman). The respective numbers for ESEC Grão-Pará North were 24 species of amphibians (23 frogs, one *Gymnophiona*) and 42 species of reptiles (24 lizards, 16 snakes, one chelonian, one caiman). Only five species of frogs, nine species of lizards, five species of snakes, one species of chelonian and one species of caiman were common to both localities. Especially the number of frogs in common was low. Another remarkable fact was that Señaris *et al.* (2008) only collected one *Gymnophthalmid* lizard (an aquatic one) and 13 species of *Hylidae*, whereas in contrast to these numbers, in ESEC Grão-Pará North we collected 11 species of *Gymnophthalmids* and only two species of *Hylidae*. Both expeditions took place in the dry season, so climate does not explain the differences found. Even though differences in habitats found in each area (e.g. presence of large rivers and Indian villages in the Guyana site, both absent in the Brazilian site) can explain part of the differences observed, they are arguably also due to the use of pitfalls in only one of the sites (Brazil) and to collector bias (collectors in the Guyana team predominantly work with frogs, those in the Brazilian Acarai team predominantly with reptiles). In our opinion this comparison reinforces

the results of the Biogeographic Similarity Coefficient, and shows the weakness of RAP's that usually only obtain a relatively small proportion of the herpetofauna available. However, combining the results of both expeditions we come to a total of 45 amphibians and 60 reptiles for this cross border area, a result closer to what could be expected than that obtained by either of the expeditions.

#### DATA FROM OTHER LOCALITIES IN CNP

As mentioned above several other studies took place in CNP, most of them not published. These studies yielded collections (in MPEG and RMNH) which provide further data for an inventory of CNP.

The studies in Jari (Monte Dourado) conducted by the University of East Anglia (Gardner *et al.*, 2007; Ribeiro-Junior *et al.*, 2008) were undertaken with a special purpose and focused on leaf litter frogs and lizards (Stokstad, 2008). Consequently, in those collections (deposited in MPEG) hardly any Hylids are present. From Jari a number of species (20) were reported that were not collected during our recent work in the CNP localities [*Bufo granulatus* Spix, 1824; *Hyla raniceps* Cope, 1862; *Scinax* sp. n. 2 (to be described by MSH shortly) [slides Enrico Bernard]; *Leptodactylus macrosternum* Miranda-Ribeiro, 1926; *Gonatodes* sp. n. (to be described shortly by Sturaro & Avila-Pires) [MPEG 23822-27; MPEG 27719]; *Cnemidophorus cryptus* Cole & Dessauer, 1993 (all female population); *Tupinambis teguixin* (Linnaeus, 1758); *Anilius scytale* (Linnaeus, 1758); *Epicrates cenchria* (Linnaeus, 1759); *Typhlops reticulata* (Linnaeus, 1766); *Atractus snethlagae* Cunha & Nascimento, 1983; *Oxybelis aeneus* (Wagler, 1824); *O. fulgidus* (Daudin, 1803); *Oxyrhopus melanogenys* (Tschudi, 1845); *Philodryas viridissimus* (Linnaeus, 1758); *Siphlophis cervinus* (Laurenti, 1768); *Spilotes pullatus* (Linnaeus, 1758); *Micrurus psyches* (Daudin, 1803); *Bothrops brazili* Hoge, 1953; *Rhinoclemmys punctularia* (Daudin, 1801)]. These species therefore can be added to the total list of species known from CNP.

Personnel of MPEG collected a further 12 species of snakes in Jari that were not collected during the project of the University of East Anglia and of which the following seven have not been collected elsewhere in CNP: *Eunectes deschauenseei* Dunn & Conant, 1936; *Helicops leopardinus* (Schlegel, 1837); *H. polylepis* Günther, 1861; *Hydrodynastes gigas* (Herrmann, 1804); *Liophis cobellus* (Linnaeus, 1758); *Liophis lineatus* (Linnaeus, 1758); *Mastigodryas bifossatus* (Raddi, 1820). Moreover they collected material of the caecilian *Microcaecila* sp. n. (to be described shortly by Maciel & Hoogmoed) [MPEG 14596-97].

Also some small collections were made in Monte Alegre that yielded some species of squamates not yet known from elsewhere in CNP. These are *Tropidurus hispidus* Spix 1825 [MPEG 24119-22, 24170-71]; *Leptotyphlops septemstriatus* (Schneider, 1801) [MPEG 21514-15]; *Epicrates maurus* Gray, 1849 [MPEG 21507-08].

Ecological herpetological work on plateaus in Floresta Nacional Saracá-Taquera (J.F.M. Sarmiento & U. Galatti, unpublished data) provided additional species (ten) that were not collected during the Calha Norte expeditions, in Jari or in Almeirim: *Hyla marmorata* (Laurenti, 1768); *Phrynohyas resinifictrix* (Goeldi, 1907); *Leptodactylus fuscus* (Schneider, 1799); *Hemidactylus mabouia* (Moreau de Jonnés, 1818); *Amphisbaena alba* (Linnaeus, 17858); *Chironius carinatus* (Linnaeus, 1758); *Dipsas variegata* (Duméril, Bibron & Duméril, 1854); *Drymarchon corais* (Boie, 1827); *Umbrivaga pygmaea* (Cope, 1868) [MPEG 20996] (this is the first record of this species for Pará); *Micrurus spixii* Wagler, 1824. The lizard *Hemidactylus mabouia*, however, is an introduced species, present only in human altered habitats and it will not be counted as part of the local herpetofauna.

Hoogmoed and Avila-Pires in 1988 collected reptiles and amphibians on the banks of rio Nhamundá (Sítio Céu Estrelado) and on the right bank of rio Trombetas (Cruz Alta). During this work they obtained several species not obtained elsewhere in CNP: *Typhlonectes compressicauda* (Duméril & Bibron, 1841); *Cnemidophorus lemniscatus*



(Linnaeus, 1758) (with males and females); *Uracentron azureum* (Linnaeus, 1758); *Helicops hagmanni* Roux, 1910; *Peltocephalus dumerilianus* (Schweigger, 1812); *Podocnemis expansa* (Schweigger, 1812); *Podocnemis unifilis* Troschel, 1848.

Morales (2002) described *Colostethus sumtuosus* from the Trombetas river. This species was not collected during our recent or any of the other expeditions.

Avila-Pires (1995) described *Tretioscincus oriximinensis* from the village of Oriximiná on the bank of the Amazon. This species was not collected during our recent or any of the other expeditions.

França *et al.* (2006) reported two species of snakes from Monte Alegre that had not been reported from CNP before: *Mastigodryas pleei* (Duméril, Bibron & Duméril, 1854) (also collected by MSH on the frontier between Brazil and Suriname (Sipaliwini savanna) [MSH 1970-68, material in RMNH] and *Phimophis guianensis* (Troschel, 1848).

A.O. Maciel & M.S. Hoogmoed, in a paper that has been submitted to Zootaxa, report *Potamotyphlus kaupii* (Berthold, 1859) from Cachoeira Porteira, Oriximiná, Pará.

Summarizing, at the moment we are aware of three species of caecilians, eight species of frogs, seven species of lizards, one species of amphisbaenid, 29 species of snakes, four species of turtles and one caiman (Medem, 1983) that have been collected in CNP, but were not collected or observed during our recent expeditions. Adding those species to our totals for CNP, we get the following totals per group: caecilians six species, anurans 85 species, lizards 43 species, amphisbaenians four species, snakes 78 species, chelonians nine species, and caiman three species. This gives a total of 90 amphibians and 137 reptiles. These numbers, which refer to all habitats present in the entire area, are getting close to what may be expected (see Introduction). However, our knowledge about the distribution of species within CNP is still very limited and mainly based on 11 localities (Figure 2) with different intensity of collecting effort.

## DATA FROM SURINAME NEAR THE BRAZILIAN BORDER

Parker (1940) described *Ninia hudsoni* from the border between Suriname and Brazil.

Hoogmoed (1969b) described *Dendrobates azureus* from a locality in Suriname close to the Brazilian border. This species was synonymised with *D. tinctorius* by Wollenberg *et al.* (2006), but we are of the opinion that this synonymisation is not correct, and that possibly the authors have been confused by using terrarium animals with unreliable locality data and by the fact that there also is a blue variant of *D. tinctorius* (collected by us in ESEC Grão-Pará North) that is similar to, but different from *D. azureus* in pattern. The pattern of *D. azureus*, as described by Hoogmoed (1969b; Lötters *et al.*, 2007: figure 708), consists of black spots haphazardly distributed on a blue background, without any reminiscence of the basic light linear pattern on the back of *D. tinctorius* (light head spot and two dorsolateral lines converging on the sacral area and continuing as a sacral line to the cloaca, [Lötters *et al.*, 2007: figures 691-700, 701-706). This basic light pattern is still distinctly recognisable in the blue form of *D. tinctorius*, present in the western part of northern Pará (Lötters *et al.*, 2007: figure 707).

Hoogmoed & Gorzula (1979) described *Ololygon trilineata* [= *Scinax trilineatus* (Hoogmoed & Gorzula, 1979)] based on material from Venezuela, Guyana and Suriname. The Suriname paratype (RMNH 18260) from the Sipaliwini savanna was collected within 7 km of the Brazilian border.

Heyer (1994) reported a specimen of *Leptodactylus pallidirostris* Lutz, 1930 from the Sipaliwini savanna, Suriname. This name now has been synonymised with *L. validus* Garman, 1887 by Yanek *et al.* (2009).

Nussbaum & Hoogmoed (1979) described *Microcaecilia taylori* Nussbaum & Hoogmoed, 1979, from a locality in Suriname (Sipaliwini) that is less than 7 km from the border with Brazil.

Hoogmoed (1977) reported *Leptotyphlops septemstriatus* (Schneider, 1801) in Suriname from two localities close to the Brazilian border.



During fieldwork in the Sipaliwini savanna in Suriname in 1968 and 1970, MSH collected *Hydrodynastes bicinctus* (Herrmann, 1804) from within 7 km of the border with Brazil [RMNH 15965 and fieldnumber MSH 1970-127, material in RMNH].

R. A. Mittermeier in 1976 collected a series of *Pseudopaludicola boliviana* Parker, 1927 on the Sipaliwini savanna in Suriname [Museum Comparative Zoology (MCZ 92418-25)]

The species mentioned above might be expected to occur in northern Pará as well, but to our knowledge have not yet been collected there. None of these species has been included in the species counts for northern Pará.

### SPECIES ABUNDANCE

Faunal assemblages in different places may have similar composition but differ in species abundance. Comparison between areas and definition of conservation measurements therefore also should take into consideration this parameter. To obtain data on abundance, however, is quite difficult, especially in tropical rainforests where many species appear to have low densities. A coarse measure of relative abundance may be obtained by looking at numbers of registered individuals, even though these numbers tend to underrepresent the most common species (where not all individuals seen are registered) and species with seclusive habits or which occupy habitats not well surveyed. Amphibians that form breeding aggregations, some only during short times, present another difficulty in such comparisons, since their numbers are not correlated with total collecting effort. Anyway, comparing numbers of registered specimens gives an idea of the most observable species in each area, during the period of the expedition. Figures 78-79, 80-81 and 82-83 show the number of specimens per species collected at each site and for all sites together, for respectively amphibians, lizards and snakes (except for snakes from FLOTA Faro and ESEC Grão-Pará Centre, where but for one species only one specimen per species was registered). Even when only

the five most common species in each area are compared, no two sites were the same. No species, in any of the groups, appear between the five most abundant in all sites. Among amphibians, *Bufo margaritifer*, *Adenomera andreae*, *Bufo* sp. n., *Anomaloglossus baeobatrachus*, and *Leptodactylus mystaceus* are the most numerous in all areas together. Considering these five species, *A. andreae* was among the five most abundant species in five of the seven sites, *B. margaritifer* in four of the sites, and *Bufo* sp. n., *Anomaloglossus baeobatrachus* and *L. mystaceus* in only two sites. Some of the most common species in one site, like *Atelopus hoogmoedi* and *Dendrobates tinctorius*, were found in no other site at all. The five most abundant lizards were *Kentropyx calcarata*, *Leposoma guianense*, *Coleodactylus amazonicus*, *Anolis chrysolepis* and *Arthrosaura reticulata*. *Leposoma guianense* was among the five most abundant species in five sites, and was the most abundant species in two of them. *Coleodactylus amazonicus* was the most abundant lizard species in three sites, and the third most abundant in a fourth site. *Kentropyx calcarata* and *Anolis chrysolepis* appeared each in three sites among the most abundant species, while *Arthrosaura reticulata* was abundant in only one site, FLOTA Faro. The five most abundant species of snakes were *Bothrops atrox*, *Liophis reginae*, *Liophis typhlus*, *Dendrophidion dendrophis* and *Leptotyphlops albifrons*. *Bothrops atrox*, the most abundant species, was represented by 13 specimens, while of the last two species of this list only five specimens were found. As pointed out before, no species was found in more than four of the sites, and only *D. dendrophis* was sampled in four sites; the other four species were registered in three sites. Differences in the most abundant species are less likely to result from sampling bias or by chance, and are more likely to reflect real differences. However, especially for amphibians part of the differences may be due to different periods of the year, but another part reflects probably differences in the available habitats – even though for most species we do not know exactly which conditions favor them.




Figure 78. Relative abundance of amphibians at each site, represented by registered number of individuals of each species. For complete species names see Appendix 1.

(Continued)






Figure 78.

(Continued)






Figure 78.

(Continued)






Figure 78.

(Conclusion)

## OBSERVATIONS ABOUT SOME SPECIES

### Amphibians

*Atelopus hoogmoedi* is a colourful small toad that until recently was known as *Atelopus spumarius* or *A. spumarius hoogmoedi* (Frost, 2009). The distribution of *A. spumarius* was supposed to reach from the Andes to the Guianas, with a gap in between those two extremes in western Amazonia. Lötters *et al.* (2002) were of the opinion that this was a species complex and that for the Guiana population the name *A. hoogmoedi* would be available. Lötters *et al.* (2005) used the name *A. hoogmoedi* for the Guiana population. *Atelopus hoogmoedi* was described from French Guiana and is known to occur throughout the three Guianas and adjacent Brazil (Noonan & Gaucher, 2005). In Pará the species was known from one small area in Monte Dourado and from a rather undefined locality "Brazil, 30 km S of the Suriname border" (material in RMNH), with outlying populations in Tucuruí, Serra de Carajás, Itaituba and near Santarém,

all four localities in Pará south of the Amazon River. The species is also known from several localities in Amapá (Lima, 2008) and from the surroundings of Manaus, Amazonas (Lima *et al.*, 2006). The Brazilian populations mentioned here are rather uniform in dorsal pattern (dark brown with vermiculate yellow to pale greenish lines on the back), but in Amapá the colour may become dark purple, with lighter purple vermiculations in some populations. The ventral colour is variable, from bright yellow everywhere, to bright yellow with bright or faint red palms, soles and seat patch, or entirely bright purple in Amapá in some populations. The genus *Atelopus* is known for the fact that many species (all from the high Andes) are threatened by extinction (a few already are extinct) probably as a result of infection with the chytrid fungus *Batrachochytrium dendrobatidis* (=Bd), which can cause the populations to collapse in a short time. Luger *et al.* (2008) checked on the viability of populations of *A. hoogmoedi* in Suriname (Brownsberg) and in Guyana (Mabura Hill Forest Reserve) and found healthy populations




Figure 79. Relative abundance of amphibians in the seven studied sites together. For complete species names see Appendix 1.




Figure 80. Relative abundance of lizards and amphibians at each site, represented by registered number of individuals of each species. For complete species names see Appendix 1.

(Continued)






Figure 80.

(Continued)






Figure 80.

(Continued)






Figure 80.

(Conclusion)

in numbers that are comparable with “those recorded for other *Atelopus* species before catastrophic declines”. During our stay in ESEC Grão-Pará Centre we encountered a large population of *A. hoogmoedi*. We collected 31 adult individuals, both males (most) and females (no juveniles) in 15 days, all during active searching, none were found in pitfalls. In daytime males were regularly heard calling. During this field period we spent 74 person-hours collecting in forest in daytime and 40 person-hours at night, a total of 114 person-hours. *Atelopus hoogmoedi* were collected both at day (most) and at night in two relatively small areas where trails crossed a creek in terra-firme forest. The species does not occur in savanna areas. Thus our collecting rate was  $31/114 = 0.272$  *A. hoogmoedi* per person-hour spent in the field, but it should be noted that only a small part of our field time was spent in *A. hoogmoedi* habitat near creeks, although this cannot be quantified. Our research was not exclusively directed at *A. hoogmoedi*, but to an inventory

of the herpetofauna in general. Thus, the value of 0.272 calculated above should be considered as a minimum value and the real population size may be considerably larger than suggested by this number. Although not directly comparable to the data from Suriname (57 specimens in 37 days) and Guyana (202 specimens, during 393.5 transect hours [0.513 individuals per transect hour]) it is our impression that the population we encountered was comparable to, or larger than, those in Suriname and Guyana and thus appears healthy. At ESEC Grão-Pará Centre we have the special situation of a large forest island in the middle of a savanna area that probably to a large extent is isolated from the terra-firme forest surrounding the savanna. Apparently there is some gallery forest along creeks that connects it in some places to the surrounding terra-firme forest. *Atelopus hoogmoedi* is restricted to that terra-firme forest island and does not occur in savanna. Whether the species occurs in the terra-firme forest surrounding our research area or in the gallery




Figure 81. Relative abundance of lizards and amphisbaenians in the seven studied sites together. For complete species names see Appendix 1.

forests is not known as we were not able to collect there, but it seems likely. The large forest island in the savanna may form a natural refuge for *A. hoogmoedi*, isolating it to a certain degree from surrounding populations in continuous terra-firme forest. In the case of a Bd infection in the terra-firme forest surrounding the savanna, the population of *A. hoogmoedi* at collecting station ESEC Grão-Pará Centre might be naturally protected against infection, just by this isolation. Considering the increasing spread of Bd (fortunately not yet found in Amazonia) this area merits special attention for conservation purposes. In contrast to most other species of *Atelopus* (except those in Suriname, French Guiana and Amapá that have the same altitudinal range as *A. hoogmoedi*),

*A. hoogmoedi* occurs only at low elevations, from 20 m at Monte Dourado, Pará, to at the most 600 m in Carajás, Pará and 700 m on the Lely Mountains in Suriname (MSH, unpublished data). According to Ron (2005) the presence of Bd at lower elevations at the moment is not very likely because medium temperatures seem to be too high for Bd infections. Thus, we might conclude that the fact that this species occurs at low altitudes provides a certain protection against Bd infection. However, this can not be considered a safeguard against Bd infection, as the way of infection and transport of the pathogen is not yet completely known.

*Bufo margaritifera* is a large species of toad (SVL females 87 mm, SVL males 66 mm [Hoogmoed, 1986],




Figure 82. Relative abundance of snakes at each site, represented by registered number of individuals of each species. No graphs are presented for FLOTA Faro, where only *Corallus hortulanus* was represented by two specimens, all other species by one specimen each, and ESEC Grão-Pará Centre, where *Leptotyphlops albifrons* was represented by three specimens, all other species by one specimen each. For complete species names see Appendix 1.

(Continued)






Figure 82.

(Continued)






Figure 82.

(Conclusion)

but in our CNP material resp. 95 mm [CN515] and 77.5 mm [CN496]) with well developed bony cranial crests, especially in the supratympanic region, which may extend vertically or horizontally, and with neural vertebral spines protruding through the dorsal skin in females, whereas males have low cranial crests and no protruding vertebral spines. In our material from FLOTA Paru, however, there is one male (CN 1726, SVL 67 mm) that has a testis, vocal slits and copulatory warts, but externally very much resembles a female by having high cranial crests, a large jaw knob and protruding neural spines. Both sexes have a distinct bony knob on the corner of the mouth (generally smaller in males) and an oblique row of tubercles from the parotoid gland to the groin (Hoogmoed, 1986). Fouquet *et al.* (2007a), thinking they were describing a new species, actually redescribed *B. margaritifera* under the name *Rhinella martyi*, as a result of not being acquainted with the species of the *B. margaritifera* group, putting too much emphasis on molecular data, working with few specimens from a restricted area and not knowing how exactly *B.*

*margaritifera* should be defined. Neither did they bother to check whether one of the many synonyms available could be used for their "new" species. Laurenti (1768, p. 30) in his description of *Rana margaritifera* refers to two drawings in Seba (1734; pl. 71, figures 6 and 7) and for his "Var.  $\beta$ " he refers to Seba (1734: pl. 71 figure 8, which shows an aberrant specimen with five fingers on the left hand, the right hand not being visible). Thus, the two specimens represented in the drawings (Seba, 1734 pl. 71 figures 6 and 7 being a dorsal and ventral view of one specimen) thus can be considered the type series of *Rana margaritifera* Laurenti, 1768. The type locality is given as Brazil ("Habitat in Brasilia"), probably based on the names used in the text in Seba (1734). Because of the problematic situation concerning this group of toads, as demonstrated by Fouquet *et al.* (2007a) and by Haas (2004) who considered a female of *B. margaritifera* with well developed crests as a male, and a male of *B. margaritifera* with low crests as a female of another species, it seems useful to indicate the specimen depicted in Seba (1734 pl. 71, figures 6 and 7) as




Figure 83. Relative abundance of snakes in the seven studied sites together. For complete species names see Appendix 1.

the lectotype of *Rana margaritifera* Laurenti, 1768, in order to avoid further confusion. The specimen depicted in Seba (1734 pl. 71, figure 8) becomes the paralectotype of *Rana margaritifera* Laurenti, 1768. This species, mentioned by Hoogmoed (1979b) as *B. typhonius*, has a distribution that at least covers Guyana, Suriname, French Guiana and, in Brazil, Amapá and Pará north and south of the Amazon. Thus, this is not a Guiana endemic. The description of *Rhinella martyi* completely agrees with the data available for the lectotype of *B. margaritifera* and the artificial distribution provided for it, due to lack of material, completely falls within the known distribution of *B. margaritifera*. Therefore, we here synonymise *R. martyi* with *B. margaritifera*. This species generally inhabits rainforest and is an explosive breeder in temporary pools and inundated areas, where large numbers may assemble. Although this species is active in daytime on the forest floor, males form choruses and call at night sitting in shallow water or on objects floating in the water (Hoogmoed, 1990b). The other species described by Fouquet *et al.* (2007a), *Rhinella lescurei*, is a good species that was already recognised by Hoogmoed (1979b) as *Bufo* sp. "B", with an altitudinal distribution of 0-300 m, and occurring in Guyana, Suriname and French Guiana (many localities [MSH, unpublished data]). Thus it is not restricted to French Guiana as suggested by Fouquet *et al.*, (2007a). During our recent surveys in CNP we did not collect this species. This small species is easily distinguished from *B. margaritifera* by its greenish iris (golden in *B. margaritifera*), a character not mentioned by Fouquet *et al.* (2007a), by several morphological characters described by Fouquet *et al.* (2007a), and by its ecology. This is a species that is active in daytime, and of which the males call singly, separated several meters from each other, from elevated posts 50 – 300 cm above the ground (rocks, leaves, palm-fronds, lianas), always near creeks in the rainforest, during daytime. They lay their eggs in small bodies of standing water near creeks (Hoogmoed, 1990b).

*Bufo* sp. n., collected in FLOTA Faro and ESEC Grão-Pará North, is a small species of the *Bufo margaritifera*

group, with green iris and no cranial crests. The species is also known from French Guiana (Hoogmoed & Avila-Pires, 1991b), Jari (Monte Dourado, Pará) and Amapá, and will be described as a new species in a forthcoming paper by MSH. It was mentioned by Hoogmoed (1979b) as *Bufo* sp. "A".

*Cochranella* sp. It has not yet been possible to identify this species with certainty. The taxonomy of Centrolenidae in the Guianan Region is still in flux and several species from other areas in eastern Amazonia, both in the Guianan Region and south of the Amazon river, still await identification. The presence of this taxon constitutes a new record for Pará. Generally Centrolenidae are considered absent from south of the Amazon River in eastern Amazonia, but several species (still to be identified) have been collected in localities in Pará like Caxiuanã, Gunma, lower rio Xingu etc., thus changing the general idea of the distribution of this family (Señaris & Ayarzagüena, 2005).

*Hyalinobatrachium iaspidiense* was collected in ESEC Grão-Pará Centre, together with *Cochranella* sp., above a creek in a large area of terra-firme forest surrounded by savanna. Yáñez-Muñoz *et al.* (2009) synonymised *H. nouraguensis* with *H. iaspidiense* and discussed some new localities for the species in Peru and Ecuador. Moreover, they mentioned localities in Venezuela, Guyana, Suriname (collected by MSH) and French Guiana and two localities in Brazil: Presidente Figueiredo, in the Guianan part of the state of Amazonas, and the lower Cristalino River in northern Mato Grosso. The present record is the first for the state of Pará.

Centrolenid larvae were collected in all three localities in ESEC Grão-Pará, but no connection could be established between them and the two species of which adults were collected in ESEC Grão-Pará Centre. In this last area an egg mass was discovered hanging from the tip of a leaf of a bush over a creek, close to the calling stations of the males of *H. iaspidiense* and *Cochranella* sp. collected there. From MSH's experience in Suriname it is clear that the eggmass does not belong to *H. iaspidiense*, which lays its clutches on the upper surface of leaves, not at the tips. Thus, the eggmass probably belongs to *Cochranella* sp. found closeby, but there

is no certainty for that. We were not successful in raising the tadpoles much beyond hatching from the eggs.

*Allobates spumaponens* was described from Guyana by Kok & Ernst (2007) and is here reported for the first time from Brazil. It was collected in all three localities in the ESEC Grão-Pará, in REBIO Maicuru and FLOTA Paru. Thus its distribution is much wider than was known up till now and it may turn out to occur throughout the western part of the Guianan Region. Identification was made by comparison with the original description (Kok & Ernst, 2007).

*Dendrobates tinctorius* (blue variant, not to be confused with *D. azureus* from Suriname) only was collected by us in ESEC Grão-Pará North, where it was very abundant throughout the area. It is a blue and black, or a black and blue, frog in which the basic *tinctorius* pattern of a light head patch and two dorsolateral stripes converging on the sacrum and continued from there as a sacral line to the cloaca, is reflected in the blue elements. The blue may be restricted to some narrow lines, often interconnected, on a black background, or it may expand to form the background color with the black reduced to a number of larger and smaller spots, still leaving the basic pattern visible. During our visit there was much courtship activity, with specimens following each other and pressing each other to the ground. *D. tinctorius* (blue variant) at present is known from ESEC Grão-Pará North, FLONA Saracá-Taquera in CNP and from the Konashen area in southern Guyana (Señaris *et al.*, 2008). All these localities are west of the Trombetas River, and this distribution is similar to that of *Leposoma* sp. n. and *Bachia panoplia*. *D. tinctorius* from eastern CNP (Jari) are black with the basic yellow pattern known for this species.

*Epipedobates* cf. *guayanensis* and *E. hahneli*. Haddad & Martins (1994) reviewed the species of the *E. pictus* group, concentrating on Brazil, and came to the conclusion that only one species, *E. hahneli*, occurred along the Amazon and its main southern tributaries. At FLOTA Trombetas we collected syntopically two small species of *Epipedobates* of similar size (25 mm svl, both with larvae and calling males – apparently in full

breeding season in mid-May, like all other *Dendrobatids* in the area). They differed in call, in body shape, and in pattern and colour of lipstripe, lateral stripe, dorsolateral stripes, belly, colour and shape of spots in the axilla, in the inguinal region, on the thighs and on the tibia. We have identified the specimens with yellow axil, inguinal and tibia spots, a white lipstripe that starts a short distance in front of the eye, no lateral white stripe and narrow white dorsolateral stripes as *E. hahneli*, a species that also occurs in Amapá (Mazagão [MPEG 810, 6936], Oiapoque [MPEG 20381-82]), Amazonas (Balbina [MPEG5966-67], Mamirauá [MPEG 7281-83, 7286-7289, 7310-7318], Urucu [MPEG 5155], Benjamin Constant [MPEG 5394-95, 5489-90, 5505-5507, 561817668]), Pará (Paraupebas [MPEG 25076-83], Belo Monte [MPEG 22008], São Felix do Xingu [MPEG 9347-48]), southern Suriname (Sipaliwini airstrip, south bank of Coeroeni River [Posts Tigrie and Gonini]) (Hoogmoed, 1971a, b as *Dendrobates pictus*) and French Guiana (Azevedo-Ramos *et al.*, 2004; Lescure & Marty, 2000; Lima, 2008; Silverstone, 1976 [part of his *Phyllobates pictus*]). The other taxon, with red spots in axil, inguina and back of thigh (the last two spots connected by a narrow orangish stripe across the dorsal surface of the thigh), a large red spot on the back of the tibia, a white lip stripe that commences close to the tip of the snout, an irregular white lateral stripe and relatively broad golden to orange dorsolateral stripes, we provisionally have identified as *E. cf. guayanensis*, a taxon that was described as *Dendrobates pictus guayanensis* from northern Venezuelan Guiana (Heatwole *et al.*, 1965), and with whose description our specimens agree well. This taxon was not considered by Haddad & Martins (1994) in their revision of the *pictus*-group, although their description of *E. pictus* from Bolivia and SW Brazil resembles our material of *E. cf. guayanensis* very much. But the distance between the localities of *E. pictus* and *E. cf. guayanensis* seems too large to consider them as one species. Lötters *et al.* (2007) reporting *E. cf. guayanensis* (as *Ameerega pictus guayanensis*) from Venezuela and



Guyana, reached the same conclusions as we do, and suggested that this may be a good species. We can not yet eliminate the possibility we are dealing with a new species here, and further study is needed to provide a decisive answer. We have located more material of *Epipedobates* cf. *guayanensis* in the collection of MPEG: from Mazagão, Amapá [MPEG 6935], from Jari (Monte Dourado), Pará [MPEG 17495-505], and from Monte Alegre, Pará [MPEG 19747-52, 20192, 20197-20203]. Only in Mazagão, Amapá, both species were also registered syntopically. *E.* cf. *guayanensis* was much more abundant (ten specimens) in FLOTA Trombetas than *E. hahneli*, of which we only collected a single specimen in a rather open area of terra-firme forest. In REBIO Maicuru we only collected one specimen of *E. hahneli* and no *E.* cf. *guayanensis*. Unfortunately no calls could be recorded. Based on the material we examined, *E.* cf. *guayanensis* does not occur south of the Amazon River, and *E. hahneli* seems to be rare north of the Amazon, although present from Amapá west through Pará to Balbina, in Amazonas. *E. hahneli* is known from south of the Amazon, from Belo Monte (Pará) in the east to Benjamin Constant (Amazonas) in the west. The eastern end of its distribution area in Brazil falls largely outside the “predicted niche” for this species computed by Twomey & Brown (2008), showing that such models should be considered with much care.

*Hyla dentei*, described from Amapá, was reported from French Guiana by Lescure & Marty (2000) and from additional localities in Amapá by Lima (2008). The species was only collected at REBIO Maicuru and is a new record for Pará.

*Hyla gaucheri* was described from coastal French Guiana and also is known from coastal and isolated savanna areas in interior Suriname (MSH, unpublished data). The species was collected near some pools and a lake in ESEC Grão-Pará South and is here reported for the first time from Brazil. The presence of pools and the lake in this locality caused the number of Hylid frog species to be considerably higher here than in any of the other localities sampled, despite the relatively advanced season (end of rainy season).

*Phrynohyas hadroceps* was described from southern Guyana (Duellman & Hoogmoed, 1992) and since has been reported from French Guiana (Lescure & Marty, 2000). Its characteristic call is a loud, regular, “woody” sound, like a metronome, that is repeated hours at an end. This call was heard (but not recorded) in REBIO Maicuru, but no specimen could be observed or collected. On the basis of the call, we report this species as new for Brazil.

*Scinax garbei* is known from Ecuador, western Brazil, adjacent Peru, Bolivia, Colombia and Venezuela (Duellman, 1972; Frost, 2009; La Marca *et al.*, 2004). Its easternmost known locality in Brazil was Manaus, Amazonas (Lima *et al.*, 2006). The record from REBIO Maicuru extends the eastern border of distribution well into Pará, for which this is a new record.

*Scinax proboscidea* was described from Suriname and since has been reported from several other localities in the Guianas (Duellman, 1972; Lescure & Marty, 2000; MSH, unpublished data) and Amapá (Lima, 2008). The record from ESEC Grão-Pará South is the first record for Pará.

*Scinax* sp. n. is a small species of *Scinax* that was only collected in REBIO Maicuru. It does not agree with any of the known species from the area and will be described as new in a separate paper.

*Scinax* gr. *ruber* is a large species related to *S. ruber*, but certainly different from it and from *S. x-signatus* (Spix, 1824). Its correct identification still has to be checked.

*Adenomera heyeri* was recently described from French Guiana (Boistel *et al.*, 2006; Angulo *et al.*, 2006) and has not yet been reported from outside that country, although MSH (unpublished data, material in RMNH) has collected it in several places in Suriname (Lely Mountains, Patamaca, Brownsberg, Kabalebo, Mozes Creek, Van Ams Creek, 20 km N. Lucie River, Oelemari, Loë Creek, Airstrip Tafelberg) and French Guiana (Mont Mahury, Mont La Gabrielle, Dégrad des Cannes) as well. The record for ESEC Grão-Pará North is the first record for Brazil.

*Eleutherodactylus chiastonotus* was known from Suriname, French Guiana and Amapá, and its presence



in northern Pará was to be expected. The records from FLOTA Trombetas and REBIO Maicuru are the first for Pará.

*Eleutherodactylus fenestratus* is generally considered a species from south of the Amazon River, but reaching Manaus and southern Guyana (Lima *et al.*, 2006). We have collected it in ESEC Grão-Pará North, which is the first record from Pará north of the Amazon.

*Eleutherodactylus inguinalis* was described from the border of Suriname and Brazil, but had not yet been reported from northern Pará, where it was to be expected, as it was already known from several localities in Suriname (MSH unpublished data, material in RMNH), French Guiana (Lescure & Marty, 2000) and Amapá (Lima, 2008). Our records from ESEC Grão-Pará North and REBIO Maicuru are the first for Pará.

*Eleutherodactylus marmoratus* is known from the three Guianas and Amapá (MacCulloch *et al.*, 2004; Lima, 2008) and its presence in northern Pará was to be expected. The record from ESEC Grão-Pará North is the first for Pará

*Leptodactylus bolivianus* has a distribution from southern Central America to Bolivia and the Guianas, but had not yet been reported from northern Pará where it was expected to occur. It is here reported from FLOTA Trombetas, ESEC Grão-Pará South, REBIO Maicuru and FLOTA Paru. The species also has been collected in Jari (Monte Dourado), Pará.

*Leptodactylus myersi* was described from Roraima (Brazil), Suriname and French Guiana, from isolated rock outcrops and granitic inselbergs in savannas and in rainforest (Heyer, 1995). The species was collected in FLOTA Trombetas in an isolated open area with flat granitic rock outcrops and boulders and a low open vegetation, surrounded by terra-firme forest, and in ESEC Grão-Pará Centre on rock outcrops (lajedos) in a savanna and in cerradão (transitional) forest bordering the savanna. However, it should be noted that in FLOTA Trombetas two specimens of this species were collected well inside the forest, in areas with large rocks, about 100 m and 300 m respectively from the open rock area. This indicates that the species does enter forest, apparently in association with rocks, at least for some distance. Heyer

(2005) reported the species from the campos de Ariramba, near Monte Alegre. The present records nicely fill the gap between the Suriname/French Guiana localities, the southern CNP locality and the Roraima localities.

*Chiasmocleis hudsoni*, a minute fossorial species, was described from southern Guyana, close to the border with Brazil. It was reported from the neighbourhood of Manaus, Amazonas, by Lima *et al.* (2006) and is also known from southern Suriname close to the Brazilian border (MSH, unpublished data). It could be expected to occur in Pará, but had not yet been reported from there and our records are the first for the state.

*Chiasmocleis* sp. n. is a small fossorial species of a genus from which recently a number of new species have been described, but all from south of the Amazon (Caramaschi & Cruz, 2001; Peloso & Sturaro, 2008) or from the Atlantic forest. The species here referred to does not agree with any of those newly described species (*C. jimi* Caramaschi & Cruz, 2001; *C. avilapiresae* Peloso & Sturaro, 2008), or with *C. shudikarensis* Dunn, 1949, known from Guyana, Suriname, French Guiana and Amazonas, Brazil (Dunn, 1949; Lescure & Marty, 2000; Lima *et al.* 2006; MSH, unpublished data). The distribution given by Rodrigues *et al.* (2004) for *C. shudikarensis* seems to be too extensive and to include distribution areas of other species as well.

*Otophryne pyburni* was known from eastern Colombia close to the Brazilian border, southern Venezuela, Suriname (Campbell & Clarke, 1998 [no locality]; MSH, unpublished data: Tepoe [RMNH MSH fieldnumber 4017] and Kwamalasemoetoe [Slide J. de Bruin]), French Guiana and Amapá (Campbell & Clarke, 1998; Lescure & Marty, 2000). Carvalho *et al.* (2007) reported it from Parque Nacional Pico de Neblina, Amazonas State, Brazil, and MacCulloch *et al.* (2008b) mentioned it from Pará, without further specification, and without indicating it in their map. We recorded it from ESEC Grão-Pará North and we also collected it at Monte Dourado, Pará [MPEG 17605]. These are the first specific localities from Pará. Señaris & Acosta-Galvis (2004), in their distribution map

(which differs from the distribution as given in the text), show a narrow band through northern Pará connecting the known distribution areas of French Guiana/Amapá with the localities in Venezuela and E. Colombia. We interpret this map as not being based on material from Pará, but as just an interpretation of a possible distribution. In fact, based on our collecting data the species most likely occurs throughout much of northern Pará and in southern Suriname (Kwamalasemoetoe and Tepoe, MSH, unpublished data). In Guyana it is known from Kartabo, in the northern part of the country (Campbell & Clarke, 1998). A distinctive character of the species is the presence of a solid, black, heart-shaped spot around the cloaca. It is visible in Figure 4 of the description by Campbell & Clarke (1998), but they did not specifically mention it. In the ESEC Grão-Pará North a large series of tadpoles was collected, including metamorphosing specimens, which will be described elsewhere.

*Synapturanus mirandaribeiroi* is a medium-sized fossorial species living just under the mat of superficial roots in rain forest, which makes it difficult to collect without using pitfalls. It was described from southern Guyana and is known from Guyana, Suriname, French Guiana, southern Venezuela, and eastern Colombia; it was reported moreover from the neighbourhood of Manaus, Amazonas (Lima *et al.*, 2006) and from Amapá (Lima, 2008), in Brazil. The records from FLOTA Faro, FLOTA Trombetas and ESEC Grão-Pará North are the first ones for Pará.

*Pipa snethlageae* was described from Belém, Pará, in 1914 and since has only rarely been collected. Trueb & Cannatella (1986) reported 13 specimens from six localities in the Amazon basin: four in Brazil, and one each in Colombia and Peru. Recently the species was reported from French Guiana by Massemin *et al.* (2003, 2007). This species is neither new for the fauna of Brazil nor for that of Pará, but it never has been reported from northern Pará. Our specimen [CN 319] was collected with a fishing net in inundated igapó forest on the bank of rio Nhamundá, in FLOTA Faro. MSH in 2006 collected a juvenile in the

rio Mutum, Amazonas (Reserva de Desenvolvimento Sustentável Cujubim) [MSH 10111 in MPEG], also in inundated forest. Another specimen [MPEG 16939] was collected in Juruti, Pará, on the south bank of the Amazon River. This seems to be a species restricted to large rivers and lakes of the Amazonian lowlands, that just enters the Guianan Region in its southern part along large rivers and in the east (French Guiana) via the coastal marshes of Amapá, as happens with other amphibians and reptiles. Its distribution area completely falls within that of *Pipa pipa*. The species is easily separated from the other species of *Pipa* by not having a skin appendage under the snout, like *Pipa pipa*, by having only a simple tubercle-like appendage at the corners of the mouth, by having a wide and short head and by having the tips of the fingers forming a square, flat surface perpendicular to the longitudinal axis of the fingers, with a pointed tubercle sticking out of each of the tips of the square.

*Rana palmipes* has a wide distribution in Amazonia and also is present in a small isolated area in northeastern coastal Brazil (northern end of the Atlantic forest). Generally the species is associated with large bodies of water like ponds and creeks. In the study area the species strangely enough only was encountered on top of the plateau where Rio Tinto's basecamp Rio Curuá (Estanifera) is established, near small pools on the road, in 'cerradão' forest (consisting of very thin small trees, standing very close together) close to the airstrip, with the nearest larger water body (rio Curuá) being hundreds of meters away at a lower elevation as well. This is the first record of this species in Pará north of the Amazon.

*Microcaecilia unicolor* was only known with certainty from French Guiana (A. O. Maciel & M.S. Hoogmoed, unpublished data), but its occurrence in this part of the Guianan Region (FLOTA Trombetas) does not come as a surprise because, like many other Gymnophiona, this species is difficult to collect and only known from relatively few specimens and localities. Earlier reports of this species from Brazil were based on mis-identified material.



*Rhinatrema* sp. n. was found in ESEC Grão-Pará North and in Porto Trombetas (also in northern Pará). It differs from *R. bivittatum* (Guérin de Méneville, 1838) in several morphological and colour characters and shortly will be described by A.O. Maciel & M.S. Hoogmoed [type-material in MPEG].

## Reptiles

*Amapasaurus tetradactylus* was collected in FLOTA Trombetas and in ESEC Grão-Pará North (Acarai Mountains). Since its description in 1970 it had not been found again until it was collected during the 2004–2006 Tumucumaque Expeditions of Conservation International, in northwestern Amapá (Lima, 2008), close to the border with Pará. The new localities here reported for Pará suggest that the species is widely distributed in northern Pará, but it is not yet possible to say whether this is a continuous distribution, or which environmental parameters define its occurrence. In the Acarai Mountains the species was collected at a short distance from the border with Guyana, so this species may turn up in southern Guyana as well.

*Bachia panoplia*, which was only known from the surroundings of Manaus, Amazonas, and from Oriximiná, Pará, was collected in FLOTA Faro, which may indicate its distribution is limited to the southwestern part of the Guianan Region, west of the Trombetas River.

*Leposoma* sp. n. still has to be described, but already was known from the surroundings of Manaus, Amazonas (Vitt *et al.*, 2008), and seems to extend its distribution at least to the western part of northern Pará, like *B. panoplia*. It was only collected in FLOTA Faro, whereas *Leposoma guianense* occurred in all studied sites and *Leposoma percarinatum*, a parthenogenetic species, in five of the seven sites. In FLOTA Faro all three species were collected, and it would be interesting to know if and how they interact.

*Gymnophthalmus* cf. *underwoodi* is a savanna inhabitant probably with a relatively large distribution in its specific habitat in southern Guiana and possibly beyond, in Alter-do-Chão, Santarém, south of the Amazon

River. Our only specimen is a female and was captured in ESEC Grão-Pará Centre under a rock on a rock slab in savanna. Whether males are present in this taxon is unknown. Most likely this is the species that was reported by Carvalho (1997) from Campos de Ariramba and Alterdo-Chão as "*Gymnophthalmus* com cauda vermelha" [= *Gymnophthalmus* with red tail], and, fleetingly, by Vanzolini & Carvalho (1991) from "northern Pará". It differs from *Gymnophthalmus vanzoi* Carvalho, 1997 by lacking a light upper lip, having two white bands bordered by black on the mentals, indistinct dorsolateral stripes, black flanks, and by having a reddish tail. It is similar to *G. underwoodi* in pattern, but differs from it by its reddish tail and by having all scales of the tail, from base to tip smooth, whereas in *G. underwoodi* the scales towards the tip of the tail have low keels, forming ten longitudinal ridges. We need to compare this specimen more extensively with material from other species/populations.

In FLOTA Paru a single specimen of *Neusticurus* was observed swimming in a creek in rain forest close to a waterfall. The specimen could not be captured, and specific identification was not possible because of light conditions. Considering what is known about the distribution of this genus in Guiana (Avila-Pires, 1995; Hoogmoed, 1973) it probably was either *N. bicarinatus* or *N. rudis*. Both species were collected in the same creek in ESEC Grão-Pará North.

*Ptychoglossus brevifrontalis* was considered a western Amazonian species, until one specimen was reported from the border of Suriname and Brazil by Hoogmoed (1973). In recent years however the species has been found in many localities in eastern and central Amazonia (Pinto & Quatman, 2005; Peloso & Avila-Pires, in press). The material from FLOTA Faro and ESEC Grão-Pará North has been incorporated in the paper by Peloso & Avila-Pires (in press) and is the first reported from Pará, but we are aware of material from several other localities in Pará south of the Amazon as well.

The small amphisbaenian *Mesobaena rhachicephalus* Hoogmoed, Pinto Rocha & Pereira, 2009, with conical, pointed snout, belongs to a genus that was only known

from southwestern Venezuela and adjacent eastern Colombia, at the edge of the Guianan Region (Gans, 1971). Its occurrence in FLOTA Faro (with two additional specimens from Porto Trombetas, Pará) (Hoogmoed *et al.*, 2009) came as a surprise. Being a fossorial animal its collecting is highly dependent on chance, but it is quite possible that its distribution is restricted to part of the Guianan Region. A number of small amphisbaenians in Amazonia show relatively small distributions (Hoogmoed & Avila-Pires, 1991a), although a closer study of them may show some of them to be synonymous with others (e.g. Hoogmoed & Mott, 2003). We can state here already that the large sample of small amphisbaenids in ESEC Grão-Pará South (19 specimens) has enabled us to establish that *Amphisbaena tragorhectes* Vanzolini, 1971, described from this area, is a junior synonym of *Amphisbaena vanzolinii*. Further arguments for this synonymisation will be provided in a forthcoming paper.

We use the name *Leptotyphlops albifrons* (Wagler, 1824) for the species that by some authors is still named *L. tenella* or *tenellus* Klauber, 1939. We consider this species to occur from Trinidad to the Guianas, but not south of the Amazon. We do not agree with the reasoning of Franco & Pinto (2009) that the name *Stenostoma albifrons* Wagler, 1824 would be a nomen dubium because of a lot of wrong identifications (which is true). In our opinion the drawing presented by Wagler (1824) clearly shows what later was described as *L. tenella* (large eyes, dark body with light zigzag lines and yellow spots on snout and tip of tail), in which we also concur with Franco & Pinto (2009). However, we think these authors are too much fixed on the type-locality given by Wagler (1824) (environs of Belém), where the species never again has been found in nearly 200 years. Spix's localities are not always reliable, and the type specimen may have been collected on the Guiana side of the Amazon. Thus there is no reason to declare *S. albifrons* Wagler a nomen dubium, but instead it becomes a senior synonym of *L. tenella* Klauber. Other material from south of the Amazon apparently has been erroneously identified.

*Leptotyphlops cupinensis* CN902A, from Grão-Pará South, was regurgitated by a half-grown *Apostolepis quinquelineatus* after it had been collected. It only concerned the posterior part of the body and tail. CN767 was collected while digging in black earth at the edge of a lake. It was inadvertently cut in several pieces, one of which was the posterior part of the body and tail, but the head could not be recovered. When dug up it was patternless bright orange. Comparison of both specimens shows them to belong to the same species – same number of scales under the tail (16), 14 scales around the middle of the tail and around the posterior part of the body, the spine at end of tail not very distinct, but present, and same body colour (no pattern), although that of CN902A was largely faded. We compared the remains of our specimens with specimens of several species of *Leptotyphlops*, *Typhlops squamosus* and *Liotyphlops ternetzii* in the MPEG collection. *T. squamosus* has a blackish dorsal region and for that reason does not qualify. *Liotyphlops ternetzii* has a high number of scales around the posterior end of the body (Dixon & Kofron, 1983), and therefore does not fit our specimens. The species of *Leptotyphlops* examined or known to us from previous studies (Hoogmoed, 1977), either have a blackish body with light zigzag stripes and a white spot on tip of tail (*L. albifrons*), a brown body with a white spot on the postanal scales (*L. collaris* Hoogmoed, 1977), a brown back and white belly (*L. dimidiatus* (Jan, 1861)), dorsal scales with brown spots (*L. macrolepis* (Peters, 1857)) or a distinct pattern of longitudinal lines (*L. septemstriatus*). Moreover, they all have either ten or 12 scales around the middle of the tail. *Leptotyphlops cupinensis* is the only *Leptotyphlops* known from the Guianan Region that is patternless and light and has 14 scales around the middle of the tail. The number of subcaudals of our two specimens falls well within the variation known for *L. cupinensis* (14-17) (Bailey & Carvalho, 1946; Orejas Miranda, 1967; Hoogmoed, 1977). We therefore came to the conclusion that the remains we have fit well with the same parts of *Leptotyphlops cupinensis* of which the colour was described by Bailey & Carvalho (1946) as “pale flesh, in alcohol creamy white, with no trace of a pigmented pattern” and by Hoogmoed



(1977), from a specimen in alcohol, as "pale yellowish brown without apparent pattern". Two slides of a live specimen of *L. cupinensis* from the rio Teles Pires, Mato Grosso, Brazil, show a similar colour as CN767 when it was dug up. It seems that *Leptotyphlops cupinensis* is the only species of *Leptotyphlops* known from the Guiana Shield that in life might be patternless bright orange. Thus, by a process of elimination and on the basis of colour, scales around posterior body, scales around the middle of the tail and number of subcaudals, we deduce that the remains we have belong to *L. cupinensis*. We realize there is margin for error, but nevertheless we are rather confident about this identification. This is the first mention of this species from Pará. From the Guianan Region it has been reported from Serra do Navio, Amapá, Brasil, and Lely Mountains, Suriname (Hoogmoed, 1977). It is not known from French Guiana.

*Typhlophis squamosus* was known from Venezuela, Guyana, Suriname and French Guiana (Kok & Rivas Fuenmayor, 2008). Cunha & Nascimento (1978) reported the species from eastern Pará, but not from north of the Amazon River. We collected a specimen in ESEC Grão-Pará Centre, but unfortunately it escaped before it could be photographed or preserved. Its identification does not pose a problem, it had the typically dark body and the light pink head known for this species (see Starace, 1998 for a picture). It is a new record for northern Pará.

*Corallus caninus* was considered a species with a wide Amazonian distribution. Recently Henderson *et al.* (2009) demonstrated that two species occurred in the Amazon area, viz. *Corallus caninus* in the Guianan region and *C. batesii* (Gray, 1860) in the rest of Amazonia. According to these authors *C. batesii* also could be a species complex. In Appendix 1 we indicate *C. caninus* therefore as a Guianan endemic.

*Apostolepis nigrolineata* generally is considered a species from south of the Amazon, where it is widely distributed (Lema & Renner, 1998; Lema, 2001). We collected it in ESEC Grão-Pará North and Centre. In the ESEC Grão-Pará North it was sympatric with *A. quinquelineata*.

*Atractus badius*. This species has been cited for a large area in Amazonia, but Hoogmoed (1980) pointed out that

it was restricted to the Guianan Region and that specimens examined from outside that region belonged to different species. He reported a specimen from Serra do Navio, Amapá. One specimen was collected in ESEC Grão-Pará South, which constitutes the first record for Pará.

*Taeniophallus nicagus* was resurrected as a valid species by Myers & Cadle (1994) and at that time was only known from Suriname. Martins & Oliveira (1998) reported it from the surroundings of Manaus. Hoogmoed and M. A. Ribeiro-Junior in 2006 collected a specimen in southern Amapá, near Mazagão, in terra-firme forest [MPEG 23312]. We collected it in ESEC Grão-Pará North and this constitutes the first record for Pará. In this locality the species was collected sympatrically with *T. brevirostris*.

*Thalesius viridis* was known only from Suriname and French Guiana (Hoogmoed, 1985; Ferreira-Yuki, 1993) until it was reported (as *Xenodon weneri* Eiselt, 1963) by Lima (2008) from the Tumucumaque Mountains, in Amapá. One specimen was collected in ESEC Grão-Pará North, and this constitutes the first record for Pará. Again, as it was collected close to the border with Guyana, it may turn up in the southern part of that country as well.

*Micrurus averyi* was described from the border of Suriname and Brazil and was already known from Manaus, Amazonas, but the record from ESEC Grão-Pará North is the first record of this species from Pará.

## CONCLUSIONS

The goal of the expeditions to northern Pará was to obtain a good impression of the herpetofauna present in the area. With a total of 80 amphibians and 95 reptiles collected or observed out of an expected total of 109 amphibians and 164 reptiles, we may conclude that the results of the expeditions were satisfactory. Taking into account material collected in other areas of northern Pará, reported in literature or present in the collection of MPEG, we even get a better result: 89 amphibians and 138 reptiles. We collected six species new to science (three frogs [*Bufo* sp. n., *Scinax* sp. n., *Chiasmocleis* sp. n.], one caecilian [*Rhinatrema* sp. n.], one



lizard [*Leposoma* sp. n., already reported in the literature], and one amphisbaenian [*Mesobaena rhachicephala*]; one species of lizard possibly new to science [*Gymnophthalmus* cf. *underwoodi*, possibly already reported in the literature]; six new records for Brazil (five frogs [*Allobates spumaponens*, *Epipedobates* cf. *guayanensis*, *Hyla gaucheri*, *Phrynohyas hadroceps*, *Adenomera heyeri*], one caecilian [*Microcaecilia unicolor*]) and 23 new records for (northern) Pará (13 frogs [*Cochranella* sp., *Hyalinobatrachium iaspidiense*, *Hyla dentei*, *Scinax garbei*, *Scinax proboscideus*, *Eleutherodactylus chiastonotus*, *E. fenestratus*, *E. inguinalis*, *E. marmoratus*, *Leptodactylus bolivianus*, *Chiasmocleis hudsoni*, *Synapturanus mirandaribeiroi*, *Rana palmipes*], four lizards [*Amapasaurus tetradactylus*, *Bachia panoplia*, *Leposoma* sp. n., *Ptychoglossus brevifrontalis*], six snakes [*Leptotyphlops cupinensis*, *Apostolepis nigrolineatus*, *Atractus badius*, *Taeniophallus nicagus*, *Thalesius viridis*, *Micrurus averyi*]). These data show that our knowledge of the herpetofauna of northern Pará has increased considerably, and has come to a level comparable to our knowledge about the herpetofauna of the two neighbouring countries Suriname and French Guiana.

It will be clear that these results are just a first step towards a better knowledge of the herpetofauna of northern Pará. We still have to learn a lot about geographic distribution within the area, and about ecological and topographic factors determining that distribution. It is hoped that the establishment of the protected areas in northern Pará will lead to a further intensification of research in that area, in order to be able to better protect the herpetofauna that still has many novelties to offer.

## FINAL REMARKS

This study of seven sites in the state of Pará, Brazil, north of the Amazon, as part of a large project aiming to establish management plans for a number of state conservation units, has allowed us to greatly improve our knowledge of this region, until now only poorly studied. These data will certainly give a better basis for establishing conservation policies for the area. However, it is important to keep in

mind that for such a large area the results obtained are just partial, and that faunistic studies in conservation units should not be limited to those that are necessary for elaborating initial management plans. It is important that long term inventories are planned and executed in conservation units, in order to effectively know their biodiversity and monitor it, and if necessary to adapt management plans to new data. Even though the use of some statistical tests, as estimators of richness (e.g., Chao 1, 2, Jackknife 1, 2, Bootstrap) may help in some analyses, it should be realized that these statistical tests are just that and they only can provide an estimate based on the data assembled, frequently over a short period. These tests do not take into account important biological (and other) factors that are of utmost importance to the organisms studied and that directly influence any estimation based on numbers collected. At best these tests can give some estimate based on the data available, just for the short period and for the specific area when and where they were obtained, and they should not be used to extrapolate data for larger areas. In the case of the herpetofauna, future studies should consider the effects of seasonality, since not all species are active throughout the year, or their apparent abundance in different periods of the year may vary. Moreover, our study indicates that many species are not evenly distributed in the whole region, but in most cases we do not know which environmental parameters are important for their distribution. Finally, not all microhabitats can be sampled adequately, and thus a number of habitat specialists (canopy, fossorial, aquatic) usually remains underrepresented. Only long term, careful studies can effectively lead to a thorough knowledge of the environment, giving better support for their conservation.

A positive development is that the northern part of Pará, together with Amapá and the neighbouring Guianan countries (Figure 1), at present form a carefully planned corridor of protected areas, following a landscape-scale approach to conservation – a core of Indian territories and more restrictive conservation units ('Estação Ecológica',

'Reserva Biológica', 'Parque Nacional', according to the Brazilian system of conservation units), surrounded by areas where the use of biodiversity is regulated to guarantee its sustainable use ('Floresta Estadual', 'Floresta Nacional', 'Reserva de Desenvolvimento Sustentável'). This gives hope that the fauna and flora of this region may escape from the threat of extinction, if indeed economic greed does not override our efforts of truly searching for a sustainable world.

## ACKNOWLEDGEMENTS

We thank all students that participated in the fieldwork: Paula Carolina de Almeida, Annelise d'Angiolella, Adriano Oliveira Maciel, Pedro Luiz Vieira del Peloso, and also M.Sc. Marco Antonio Ribeiro Junior, all of whom had to work hard, often under adverse conditions, but at the same time learned the pleasant and not-so-pleasant aspects of fieldwork in isolated places. Weverton Souza Bandeira Mota was part of the ichthyological team, but in his spare moments at night joined the herpetological team as much as possible, thus effectively increasing its efficiency. Aduauto Lima Cardoso tried to compensate the poor collecting results of mammal traps by joining the herpetological team at night and provided us with some interesting specimens. A special word of thanks is due to the field assistants that installed the pitfalls and drift fences and also participated in checking the pitfalls, at times installed at places that required long walks in hilly terrain: Zailson Santos da Silva, Francisco Alves Martins and Ronei de Sousa Oliveira. Also many other participants in the expeditions contributed to the final inventory by contributing specimens, or photos that served as records for the presence of certain species, even if not collected. Ariane Araujo provided slides of live *Leptotyphlops cupinensis* that were made by Robson W. Ávila. Luis Barbosa of Conservação Internacional (Belém) provided the maps.

Conservação Internacional (CI-Brasil) financed travel from Belém to Santarém and Boa Vista, the costs of the boat for the first expedition, subsistence costs of all expeditions, as well as grants to some of the researchers. Waldima Alves da Rocha wants to express her gratitude

to CI-Brasil for the grant that enabled her to participate in the project and in publishing the results. As conditioned by the mining research licence given by SEMA-PA, mining company Rio Tinto, exploring for bauxite in the area, covered all costs regarding logistics from Santarém and Boa Vista to the campsites, and took care of opening up heliports, mounting and disassembling camps, flying in personal, food and material, and removing all extraneous material from the campsites. Personnel of SEMA-Pará gave us all support within their possibilities.

## REFERENCES

- ANGULO, A., J.-C. MASSARY & R. BOISTEL, 2006. Addenda and errata to "Description of a new species of the genus *Adenomera* (Amphibia, Anura, Leptodactylidae) from French Guiana". **Acta Herpetologica** 1(2): 159-162.
- AVILA-PIRES, T. C. S., 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). **Zoologische Verhandlungen Leiden** 299: 1- 706.
- AVILA-PIRES, T. C. S., 2005. Reptiles. In: T. HOLLOWELL & R. P. REYNOLDS (Eds.): Checklist of the terrestrial vertebrates of the Guiana Shield. **Bulletin of the Biological Society of Washington** 13: 22-40.
- AVILA-PIRES, T. C. S., M. S. HOOGMOED & L. J. VITT, 2007. Herpetofauna da Amazônia. In: L. B. NASCIMENTO & M. E. OLIVEIRA (Eds.): **Herpetologia no Brasil II**: 13-43. Sociedade Brasileira de Herpetologia, Belo Horizonte.
- AVILA-PIRES, T. C. S., L. J. VITT, S. S. SARTORIUS & P. A. ZANI, 2009. Squamate (Reptilia) from four sites in southern Amazonia, with a biogeographic analysis of Amazonian lizards. **Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais** 4(2): 99-118.
- AZEVEDO-RAMOS, C., L. A. COLOMA, S. RON, F. CASTRO & C. GASCON, 2004. *Ameerega hahneli*. In: **IUCN 2009. IUCN Red List of Threatened Species. Version 2009.1**. Disponible in: <www.iucnredlist.org>. Accessed on: 23 August 2009.
- BAILEY, J. R. & A. L. CARVALHO, 1946. A new *Leptotyphlops* from Mato Grosso, with notes on *Leptotyphlops tenella* Klauber. **Boletim do Museu Nacional. Nova Série Zoologia** 52: 1-7.
- BARTLETT, R. D. & P. BARTLETT, 2003. **Reptiles and amphibians of the Amazon. An ecotourist guide**: 1-291. University Press of Florida, Gainesville.
- BOISTEL, R., J.-C. MASSARY & A. ANGULO, 2006. Description of a new species of the genus *Adenomera* (Amphibia, Anura, Leptodactylidae) from French Guiana. **Acta Herpetologica** 1(1): 1-14.



- CALDWELL, J. P. & M. S. HOOGLMOED, 1998. Allophryniidae, *Allophryne*, *A. ruthveni*. **Catalogue of American Amphibians and Reptiles** 666: 1-3.
- CAMPBELL, J. A. & B. T. CLARKE, 1998. A review of frogs of the genus *Otophryne* (Microhylidae) with the description of a new species. **Herpetologica** 54(3): 301-317.
- CAMPBELL, J. A. & W. W. LAMAR, 2004. **The venomous reptiles of the western hemisphere**: i-xviii, 1-476, [1-28], i-xiv, 477-870, [1-28]. Cornell University Press, Ithaca and London.
- CARAMASCHI, U. & C. A. G. CRUZ, 2001. A new species of *Chiasmocleis* from Brazilian Amazônia (Amphibia, Anura, Microhylidae). **Boletim do Museu Nacional, Nova Série Zoologia** 469: 1-8.
- CARVALHO, C. M., 1997. Uma nova espécie de Microteiideo do gênero *Gymnophthalmus* do estado de Roraima, Brasil (Sauria, Gymnophthalmidae). **Papéis Avulsos de Zoologia** 40(10): 161-174.
- CARVALHO, C. M., 2002. Descrição de uma nova espécie de *Micrurus* do estado de Roraima, Brasil (Serpentes, Elapidae). **Papéis Avulsos de Zoologia** 42(8): 183-192.
- CARVALHO, V. T.; L. BONORA & R. C. VOGT, 2007. Geographic distribution: *Otophryne pyburni*. **Herpetological Review** 38(3): 349.
- CECHIN, S. Z. & M. MARTINS, 2000. Eficiência de armadilhas de queda (pitfall traps) em anfíbios e répteis no Brasil. **Revista Brasileira de Zoologia** 117(3): 729-740.
- CHIPPEAUX, J. P., 1986. **Les serpents de la Guyane française**: 1-165. Éditions de l'ORSTOM (Faune Tropicale, 27), Paris.
- COLE, C. J. & P. J. R. KOK, 2006. A new species of gekkonid lizard (Sphaerodactylinae: *Gonatodes*) from Guyana, South America. **American Museum Novitates** 3524: 1-13.
- COLWELL, R. K., 2005. **EstimateS: Statistical estimation of species richness and shared species from samples**. Version 7.5. Sinauer Associates, Sunderland, Massachusetts.
- CORN, P. S., 1994. Straight line drift fences and pitfall traps. In: W. R. HEYER, M. A. DONNELLY, R. W. MCDIARMID, L. C. HAYEK & M. S. FOSTER (Eds.): **Measuring and monitoring biological diversity. Standard methods for amphibians**: 109-117. Smithsonian Institution Press, Washington.
- CRUMP, M. L. & N. J. SCOTT, 1994. Visual encounter surveys. In: W. R. HEYER, M. A. DONNELLY, R. W. MCDIARMID, L. C. HAYEK & M. S. FOSTER (Eds.): **Measuring and monitoring biological diversity. Standard methods for amphibians**: 84-92. Smithsonian Institution Press, Washington.
- CUNHA, O. R. & F. P. NASCIMENTO, 1978. Ofídios da Amazônia. X. As cobras da região leste do Pará. **Publicações Avulsas do Museu Paraense Emílio Goeldi** 32: 1-218.
- CUNHA, O. R., F. P. NASCIMENTO & A. R. HOGE, 1980. Ofídios da Amazônia. XI. Ofídios de Roraima e notas sobre *Erythrolamprus bauperthuisii* (Linnaeus, 1752). **Boletim do Museu Paraense Emílio Goeldi, série Zoologia** 120: 1-21.
- DIXON, J. R. & C. P. KOFRON, 1983. The Central and South American Anomalepid snakes of the genus *Liotyphlops*. **Amphibia-Reptilia** 4(2-4): 241-264.
- DIXON, J. R., J. A. WIEST JR. & J. M. CEI, 1993. Revision of the neotropical snake genus *Chironius* Fitzinger (Serpentes, Colubridae). **Museu Regionale di Scienze Naturali Monografie** 13: 1-279.
- DONNELLY, M., M. CHEN, C. WATSON & G. G. WATKINS, 1998. Amphibians and Reptiles of the Iwokrama forest. In: ANONYMOUS (Ed.) **The vertebrate fauna of the Iwokrama Forest. Final report from work carried out in the Iwokrama Forest by the Academy of Natural Sciences of Philadelphia 1996-1998**: 18-36.
- DONNELLY, M. A., M. H. CHEN & G. G. WATKINS, 2005a. Sampling amphibians and reptiles in the Iwokrama Forest ecosystem. **Proceedings of the Academy of Natural Sciences of Philadelphia** 154: 55-69.
- DONNELLY, M. A., M. H. CHEN & G. G. WATKINS, 2005b. The Iwokrama herpetofauna: an exploration of diversity in a Guyanan rainforest. In: M. A. DONNELLY, B. I. CROTHER, C. GUYER, M. H. WAKE & M. E. WHITE (Eds.): **Ecology and evolution in the tropics: a herpetological perspective**: 428-460. University of Chicago Press, Chicago.
- DONNELLY, M. A., R. D. MACCULLOCH, C. A. UGARTE & D. KIZIRIAN, 2006. A new riparian Gymnophthalmid (Squamata) from Guyana. **Copeia** 2006(3): 396-403.
- DUELLMAN, W. E., 1972. South American frogs of the *Hyla rostrata* group (Amphibia, Anura, Hylidae). **Zoologische Mededelingen Leiden** 47: 177-192.
- DUELLMAN, W. E., 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. **Miscellaneous Publications, Museum of Natural History, University of Kansas** 65: 1-352.
- DUELLMAN, W. E., 1990. Herpetofaunas in Neotropical rainforests: comparative composition, history, and resource use. In: A. H. GENTRY (Ed.): **Four Neotropical Rainforests**: 455-505. Yale University Press, New Haven.
- DUELLMAN, W. E., 1999. Distribution patterns of amphibians in South America. In: W. E. DUELLMAN (Ed.): **Patterns of distribution of amphibians: a global perspective**: 255-328. John Hopkins University Press, Baltimore.
- DUELLMAN, W. E., 2005. **Cusco Amazónico. The lives of amphibians and reptiles in an Amazonian forest**: i-xv, 1-433. Comstock Publishing Association, Ithaca and London.

DUELLMAN, W. E. & M. S. HOOGMOED, 1992. Some hylid frogs from the Guiana highlands, northeastern South America: new species, distributional records, and a generic reallocation. **Occasional Papers of the Museum of Natural History, University of Kansas** 147: 1-21.

DUNN, E. R., 1949. Notes on South American frogs of the family Microhylidae. **American Museum Novitates** 1419: 1-21.

EMC (Environmental Management Consultants), 2006. **Rapid biodiversity assessment of Halcrow and Guysuco conservancies (E1751). Final Report:** 1-47. Disponível in: <[http://www-wds.worldbank.org/external/default/WDSCContentServer/WDSP/IB/2007/11/12/000020439\\_20071112132643/Rendered/PDF/E17510Final0Report010GUYSUUCOORBA.pdf](http://www-wds.worldbank.org/external/default/WDSCContentServer/WDSP/IB/2007/11/12/000020439_20071112132643/Rendered/PDF/E17510Final0Report010GUYSUUCOORBA.pdf)>. Accessed on: 12 March 2010.

ERNST, R., M.-O. RÖDEL & D. ARJOON, 2005. On the cutting edge – The anuran fauna of the Mabura Hill Forest Reserve, Central Guyana. **Salamandra** 41(4): 179-194.

ERNST, R., T. KONRAD, K. E. LINSENMAIR & M.-O. RÖDEL, 2007. Impacts of selective logging on three sympatric species of *Leptodactylus* in a Central Guyana rainforest. **Amphibia-Reptilia** 28: 51-64.

EVA, H. D. & O. HUBER (Eds.), 2005. **EUR 21808 A proposal for defining the geographical boundaries of Amazonia.** Synthesis of the results from an Expert Consultation Workshop organized by the European Commission in collaboration with the Amazon Cooperation Treaty Organization: i-x + 1-39. Office for Official Publications of the European Communities, Luxembourg.

FAIVOVITCH, J., C. F. B. HADDAD, P. C. D. A. GARCIA, D. R. FROST, J. A. CAMPBELL & W. C. WHEELER, 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: a phylogenetic analysis and taxonomic revision. **Bulletin of the American Museum of Natural History** 294: 1-240.

FERREIRA-YUKI, V. L., 1993. Realocação genérica de *Xenodon werneri* Eiselt, 1963 (Serpentes: Colubridae). **Comunicações do Museu de Ciências da PUCRS** 50-54: 39-47.

FOUQUET, A., P. GAUCHER, M. BLANC & C. M. VELEZ-RODRIGUES, 2007a. Description of two new species of *Rhinella* (Anura: Bufonidae) from the lowlands of the Guiana Shield. **Zootaxa** 1663: 17-32.

FOUQUET, A., M. VENCES, M.-D. SALDUCCI, A. MEYER, C. MARTY, M. BLANC & A. GILLES, 2007b. Revealing cryptic diversity using molecular phylogenetics and phylogeography in frogs of the *Scinax ruber* and *Rhinella margaritifera* species groups. **Molecular phylogenetics and Evolution** 45: 567-582.

FRANÇA, F. G. R., D. O. MESQUITA & G. R. COLLI, 2006. A checklist of snakes from Amazonian savannas in Brazil, housed in the Coleção Herpetológica da Universidade de Brasília, with new distribution records. **Occasional Papers Sam Noble Oklahoma Museum of Natural History** 17: 1-13.

FRANCO, F. L., M. G. SALOMÃO & P. AURICCHIO, 2002. Répteis. In: P. AURICCHIO & M. G. SALOMÃO (Eds.): **Manual de Técnicas e Preparação de Vertebrados:** 77-121. Instituto Pau-Brasil de História Natural, São Paulo.

FRANCO, F. L. & R. R. PINTO, 2009. *Stenostoma albifrons* Wagler in Spix, 1824 as nomen dubium and recognition of the name *Leptotyphlops tenellus* Klauber, 1939 (Serpentes: Leptotyphlopidae). **Salamandra** 45(4): 239-244.

FROST, D. R., 2009. **Amphibian species of the world: an online reference.** American Museum of Natural History, New York. Version 5.3. Disponível in: <<http://research.amnh.org/herpetology/amphibia/index.php>>. Access on: 12 February 2009.

FROST, D. R., T. GRANT, J. FAIVOVITCH, R. H. BAIN, A. HAAS, C. F. B. HADDAD, R. O. DE SÁ, A. CHANNING, M. WILKINSON, S. C. DONNELLAN, C. J. RAXWORTHY, J. A. CAMPBELL, B. L. BLOTTO, P. E. MOLER, R. C. DREWES, R. A. NUSSBAUM, J. D. LYNCH, D. M. GREEN & W. C. WHEELER, 2006. The amphibian tree of life. **Bulletin of the American Museum of Natural History** 297: 1-370.

GANS, C., 1971. Redescription of three monotypic genera of amphisbaenians from South America: *Aulura* Barbour, *Bronia* Gray, and *Mesobaena* Mertens. **American Museum Novitates** 2475: 1-32.

GARDNER, T. A., M. A. RIBEIRO-JUNIOR, J. BARLOW, T. C. S. AVILA-PIRES, M. S. HOOGMOED & C. A. PERES, 2007. The value of primary, secondary, and plantation forests for a neotropical herpetofauna. **Conservation Biology** 21(3): 775-787.

GASC, J.-P., 1990. **Les lézards de Guyane:** 1-76. Ed. Raymond Chabanaud, Paris.

GASC, J.-P. & M. T. RODRIGUES, 1980. Liste préliminaire des serpents de la Guyane française. **Bulletin Museum National d'Histoire Naturelle** (4) 2A(2): 559-598.

GORZULA, S. & J. C. SEÑARIS, 1999. Contribution to the herpetofauna of the Venezuelan Guayana. I. A data base. **Scientia Guianae** 8: i-xviii, 1-269.

GOVERNO DO ESTADO DO PARÁ, 2006. **Pará cria maior área contínua de preservação do mundo.** Disponível in: <<http://www.amazonia.org.br/noticias/noticia.cfm?id=227652>>. Accessed on: 4 February 2010.

GRANT, T., D. R. FROST, J. P. CALDWELL, R. GAGLIARDO, C. F. B. HADDAD, P. J. R. KOK, D. B. MEANS, B. P. NOONAN, W. E. SCHARGEL & W. C. WHEELER, 2006. Phylogenetic systematics of dart-poison frogs and their relatives (Amphibia: Athespatanura: Dendrobatidae). **Bulletin of the American Museum of Natural History** 299: 1-262.

HAAS, W., 2004. Beitrag zur Biologie und Verbreitung dreier Vertreter des neotropischen *Bufo typhonius*-Komplexes (Anura: Bufonidae). **Salamandra** 40(3/4): 207-216.



- HADDAD, C. F. B. & M. MARTINS, 1994. Four species of Brazilian poison frogs related to *Epipedobates pictus* (Dendrobatidae): taxonomy and natural history observations. **Herpetologica** 59(3): 282-295.
- HALLER, E. C. P. & M. T. RODRIGUES, 2005. *Podocnemis unifilis* (Yellow-spotted River Turtle). Nests and Nesting. **Herpetological Review** 36(1): 60.
- HALLER, E. C. P. & M. T. RODRIGUES, 2006. Reproductive biology of the Amazonian Turtle *Podocnemis sextuberculata* (Testudinata: Pelomedusidae) in the Biological Reserve of Rio Trombetas, Pará, Brazil. **Chelonian Conservation and Biology** 5: 280-284.
- HEATWOLE, H., H. SOLANO & A. HEATWOLE, 1965. Notes on amphibians of the Venezuelan Guayanas with description of two new forms. **Acta Biologica Venezuelica** 4(12): 349-364.
- HEDGES, S. B., W. E. DUELLMAN & M. P. HEINICKE, 2008. New world direct-developing frogs (Anura: Terrarana): molecular phylogeny, classification, biogeography, and conservation. **Zootaxa** 1737: 1-182.
- HEINICKE, M. P., W. E. DUELLMAN & S. B. HEDGES, 2007. Major Caribbean and Central American frog faunas originated by ancient oceanic dispersal. **Proceedings of the Academy of Natural Sciences** 104: 10092-10097.
- HENDERSON, R. W., P. PASSOS & D. FEITOSA, 2009. Geographic variation in the emerald treeboa, *Corallus caninus* (Squamata: Boidae). **Copeia** 2009(3): 572-582.
- HEYER, W. R., 1994. Variation within the *Leptodactylus podicipinus-wagneri* complex of frogs (Amphibia: Leptodactylidae). **Smithsonian Contributions to Zoology** 546: 1-124.
- HEYER, W. R., 1995. South American rocky habitat *Leptodactylus* (Amphibia: Anura: Leptodactylidae) with description of two new species. **Proceedings of the Biological Society of Washington** 108(4): 695-716.
- HEYER, W. R., 2005. Variation and taxonomic clarification of the large species of the *Leptodactylus pentadactylus* species group (Amphibia: Leptodactylidae) from Middle America, northern South America, and Amazonia. **Arquivos de Zoologia** 37(3): 269-348.
- HOLLOWELL, T. & R. P. REYNOLDS (Eds.), 2005. Checklist of the terrestrial vertebrates of the Guiana Shield. **Bulletin of the Biological Society of Washington** 13: i-ix, 1-98.
- HOOGMOED, M. S., 1969a. Notes on the fauna of Surinam II. – On the occurrence of *Allophryne ruthveni* Gaige (Amphibia, Salientia, Hylidae) in Surinam. **Zoologische Mededelingen Leiden** 44(5): 75-81.
- HOOGMOED, M. S., 1969b. Notes on the herpetofauna of Surinam III. – A new species of *Dendrobates* (Amphibia Salientia, Dendrobatidae) from Surinam. **Zoologische Mededelingen Leiden** 44(9): 133-141.
- HOOGMOED, M. S., 1971a. *Dendrobates*, eine farbenreiche Gattung. **Die Aquarien- und Terrarien-Zeitschrift (DATZ)** 24(1): 1-7.
- HOOGMOED, M. S., 1971b. *Dendrobates*, een kleurrijk genus. **Het Aquarium** 41(8): 182-189.
- HOOGMOED, M. S., 1973. **Notes on the herpetofauna of Surinam**. IV. The lizards and amphisbaenians of Surinam: 1-417. Dr. W. Junk Publishers (Biogeographica, v. 4), The Hague.
- HOOGMOED, M. S., 1977. On a new species of *Leptotyphlops* from Surinam, with notes on the other Surinam species of the genus (Leptotyphlopidae, Serpentes). Notes on the herpetofauna of Surinam V. **Zoologische Mededelingen Leiden** 51(7): 99-123.
- HOOGMOED, M. S., 1979a. Resurrection of *Hyla ornatissima* Noble (Amphibia, Hylidae) and remarks on related species of green tree frogs from the Guiana area. Notes on the herpetofauna of Surinam VI. **Zoologische Verhandelingen Leiden** 172: 1-46.
- HOOGMOED, M. S., 1979b. The herpetofauna of the Guianan Region. In: W. E. DUELLMAN (Ed.): The South American herpetofauna: its origin, evolution and dispersal. **Museum of Natural History, The University of Kansas Monographs** 7: 241-279.
- HOOGMOED, M. S., 1980. Revision of the genus *Atractus* in Surinam, with the resurrection of two species (Reptilia, Colubridae). Notes on the herpetofauna of Surinam VII. **Zoologische Verhandelingen Leiden** 175: 1-47.
- HOOGMOED, M. S., 1983 [1982]. Snakes of the Guianan Region. **Memórias do Instituto Butantan** 46: 219-254.
- HOOGMOED, M. S., 1985. *Xenodon werneri* Eiselt, a poorly known snake from Guiana, with notes on *Waglerophis merremii* (Wagler) (Reptilia: Serpentes: Colubridae). Notes on the Herpetofauna of Surinam IX. **Zoologische Mededelingen Leiden** 59(8): 79-88.
- HOOGMOED, M. S., 1986. Biosystematic studies of the *Bufo* "typhonius" group. A preliminary progress report. In: Z. ROČEK (Ed.): **Studies in Herpetology**: 147-150. Societas Europaea Herpetologica, Prague.
- HOOGMOED, M. S., 1990a. Resurrection of *Hyla wavnini* Parker (Amphibia: Anura: Hylidae), a gladiator frog from northern South America. **Zoologische Mededelingen Leiden** 64(6): 71-93.
- HOOGMOED, M. S., 1990b. Biosystematics of South American Bufonidae, with special reference to the *Bufo* "typhonius" group. In: G. PETERS & R. HÜTTERER (Eds.): **Vertebrates in the tropics**: 113-123. Museum Alexander Koenig, Bonn.
- HOOGMOED, M. S. & T. C. S. AVILA-PIRES, 1989. Observations on the nocturnal activity of lizards in a marshy area in Serra do Navio, Brazil. **Tropical Zoology** 2: 165-173.
- HOOGMOED, M. S. & T. C. S. AVILA-PIRES, 1991a. A new species of *Amphisbaena* (Reptilia: Amphisbaenia: Amphisbaenidae) from western Amazonian Brazil. **Boletim do Museu Paraense Emílio Goeldi, série Zoologia** 7(1): 77-94.



- HOOGMOED, M. S. & T. C. S. AVILA-PIRES, 1991b. Annotated checklist of the herpetofauna of Petit Saut, Sinnamary River, French Guiana. **Zoologische Mededelingen Leiden** 65(5): 53-88.
- HOOGMOED, M. S. & T. C. S. AVILA-PIRES, 1992. Studies on the species of the South American lizard genus *Arthrosaura* Boulenger (Reptilia: Sauria: Teiidae), with the resurrection of two species. **Zoologische Mededelingen Leiden** 66(35): 453-484.
- HOOGMOED, M. S. & S. J. GORZULA, 1979. Checklist of the savanna inhabiting frogs of the El Manteco region with notes on their ecology and the description of a new species of tree frog (Hylidae, Anura). **Zoologische Mededelingen Leiden** 54(13): 183-216.
- HOOGMOED, M. S. & J. LESCURE, 1975. An annotated checklist of the lizards of French Guiana, mainly based on two recent collections. **Zoologische Mededelingen Leiden** 49(13): 141-171.
- HOOGMOED, M. S. & T. MOTT, 2003. On the identity of *Amphisbaena hugoi* Vanzolini, 1990 (Reptilia: Squamata: Amphisbaenidae). **Zoologische Mededelingen Leiden** 77(28): 455-457.
- HOOGMOED, M. S., R. R. PINTO, W. A. ROCHA & E. G. PEREIRA, 2009. A new species of *Mesobaena* Mertens, 1925 (Squamata: Amphisbaenidae) from Brazilian Guiana, with a key to the Amphisbaenidae of the Guianan Region. **Herpetologica** 65(4): 436-448.
- KOK, P. J. R., 2000. A survey of the anuran fauna of Montagne Belvédère, county of Saül, French Guiana: field list with comments on taxonomy and ecology. **The British Herpetological Society Bulletin** 71: 6-26.
- KOK, P. J. R., 2005. A new genus and species of gymnophthalmid lizard (Squamata: Gymnophthalmidae) from Kaieteur National Park, Guyana. **Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie** 75: 35-45.
- KOK, P. J. R., 2006a. A new species of *Hypsiboas* (Amphibia: Anura: Hylidae) from Kaieteur National Park, eastern edge of the Pakaraima Mountains, Guyana. **Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie** 76: 191-200.
- KOK, P. J. R., 2006b. A new snake of the genus *Atractus* Wagler, 1828 (Reptilia: Squamata: Colubridae) from Kaieteur National Park, Guyana, northeastern South America. **Zootaxa** 1378: 19-35.
- KOK, P. J. R., 2008a. A new highland species of *Arthrosaura* Boulenger, 1885 (Squamata: Gymnophthalmidae) from Maringma tepui on the border of Guyana and Brazil. **Zootaxa** 1909: 1-15.
- KOK, P. J. R., 2008b. Lizard in the clouds: a new highland genus and species of Gymnophthalmidae (Reptilia: Squamata) from Maringma tepui, western Guyana. **Zootaxa** 1992: 53-67.
- KOK, P. J. R., 2009. A new species of *Oreophrynella* (Anura: Bufonidae) from the Pantepui region of Guyana, with notes on *O. macconnelli* Boulenger, 1900. **Zootaxa** 2071: 35-49.
- KOK, P. J. R., R. D. MACCULLOCH, P. GAUCHER, E. H. POELMAN, G. R. BOURNE, A. LATHROP & G. L. LENGLET, 2006a. A new species of *Colostethus* (Anura: Dendrobatidae) from French Guiana with a redescription of *Colostethus beebei* (Noble, 1923) from its type locality. **Phyllomedusa** 5(1): 43-66.
- KOK, P. J. R., H. SAMBHU, I. ROOPSIND, G. L. LENGLET & G. R. BOURNE, 2006b. A new species of *Colostethus* (Anura: Dendrobatidae) with maternal care from Kaieteur National Park, Guyana. **Zootaxa** 1238: 35-61.
- KOK, P. J. R., M. N. C. KOKUBUM, R. D. MACCULLOCH & A. LATHROP, 2007. Morphological variation in *Leptodactylus lutzi* (Heyer, 1975) (Anura: Leptodactylidae) with description of its advertisement call and notes on its courtship behavior. **Phyllomedusa** 6(1): 45-60.
- KOK, P. J. R. & R. ERNST, 2007. A new species of *Allobates* (Anura: Aromobatidae: Allobatinae) exhibiting a novel reproductive behaviour. **Zootaxa** 1555: 21-38.
- KOK, P. J. R. & S. CASTROVIEJO-FISHER, 2008. Glassfrogs (Anura: Centrolenidae) of Kaieteur National Park, Guyana, with notes on the distribution and taxonomy of some species of the family in the Guiana Shield. **Zootaxa** 1680: 25-53.
- KOK, P. J. R. & M. KALAMANDEEN, 2008. Introduction to the taxonomy of the amphibians of Kaieteur National Park, Guyana. **Abc Taxa** 5: 1-278.
- KOK, P. J. R. & G. RIVAS FUENMAYOR, 2008. *Typhlophis ayarzaguenai* Señaris, 1998 is a junior synonym of *Typhlophis squamosus* (Schlegel, 1837). **Amphibia-Reptilia** 29(4): 555-558.
- LA MARCA, E., C. AZEVEDO-RAMOS, L. A. COLOMA & S. RON, 2004. *Scinax garbei*. In: IUCN 2009. **IUCN Red List of Threatened Species. Version 2009.1**. Disponible in: <[www.iucnredlist.org](http://www.iucnredlist.org)>. Accessed on: 23 August 2009.
- LATHROP, A. & R. D. MACCULLOCH, 2007. A new species of *Oreophrynella* (Anura: Bufonidae) from the highlands of Guyana. **Herpetologica** 63: 87-93.
- LAURENTI, J. N., 1768. *Specimen medicum, exhibens synopsis Reptilium emendatam cum experimentis circavenena et antidota reptilium austriacorum*: 1-214. Trattner, Vienna.
- LEMA, T., 2001. Fossorial snake genus *Apostolepis* from South America (Serpentes: Colubridae: Elapomorphae). **Cuadernos de Herpetologia** 15(1): 29-43.
- LEMA, T. & M. F. RENNER, 1998. O status de *Apostolepis quinquelineata* Boulenger, 1896, *A. pygmaea*, Boulenger, 1903, e *A. rondoni* Amaral, 1925 (Serpentes, Colubridae, Elapomorphi). **Biociências** 6(1): 99-121.
- LESCURE, J. & C. MARTY, 2000. Atlas des amphibiens de Guyane. **Patrimoines Naturels** 45: 1-388.



- LIMA, A. P., W. E. MAGNUSSON, M. MENIN, L. K. ERDMANN, D. J. RODRIGUES, C. KELLER & W. HÖDL, 2006. **Guia de sapos da Reserva Adolpho Ducke, Amazônia Central**: 1-168. INPA, Manaus.
- LIMA, J. D., 2008. A herpetofauna do Parque Nacional Montanhas do Tumucumaque, Amapá, Brasil, Expedições I – V. In: E. BERNARD (Ed.): *Inventários Biológicos Rápidos no Parque Nacional Montanhas do Tumucumaque, Amapá, Brasil*. **RAP Bulletin of Biological Assessment** 48: 38-50. Disponible in: <[http://www.conservation.org.br/publicacoes/files/RAP\\_Tumucumaque.pdf](http://www.conservation.org.br/publicacoes/files/RAP_Tumucumaque.pdf)>. Accessed on: 12 March 2010.
- LÖTTERS, S., W. HAAS, S. SCHICK & W. BÖHME, 2002. On the systematics of the harlequin frogs (Amphibia: Bufonidae: *Atelopus*) from Amazonia. II. Redescription of *Atelopus pulcher* (Boulenger, 1882) from the eastern Andean versant in Peru. **Salamandra** 38: 165-184.
- LÖTTERS, S., R. BOISTEL, M. BLANC, C. F. B. HADDAD & A. VAN DER MEIDEN, 2005. *Atelopus hoogmoedi*. In: J. E. RUEDA-ALMONACID, J. V. RODRIGUEZ-MAHECHA, E. LA MARCA, S. LÖTTERS, T. KAHN & A. ANGULO (Eds.): **Ranas arlequines**: 132-134. Conservacion Internacional (Serie libretas de campo, 5), Bogotá.
- LÖTTERS, S., K.-H. JUNGFER, F. W. HENKEL & W. SCHMIDT, 2007. **Poison frogs. Biology, species & captive care**: 1-668. Edition Chimaira, Frankfurt.
- LUGER, M., T. W. J. GARNER, R. ERNST, W. HÖDL & S. LÖTTERS, 2008. No evidence for precipitous declines of harlequin frogs (*Atelopus*) in the Guyanas. **Studies on Neotropical Fauna and Environment** 43(3): 177-180.
- MACCULLOCH, R. D. & A. LATHROP, 2001. A new species of *Arthrosaura* (Sauria: Teiidae) from the highlands of Guyana. **Caribbean Journal of Science** 37: 174-181.
- MACCULLOCH, R. D. & A. LATHROP, 2002. Exceptional diversity of *Stefania* (Anura: Hylidae) on Mount Ayanganna, Guyana: three new species and new distribution records. **Herpetologica** 58: 327-346.
- MACCULLOCH, R. D. & A. LATHROP, 2004a. A new species of *Dipsas* (Squamata: Colubridae) from Guyana. **Revista de Biología Tropical** 52: 239-247.
- MACCULLOCH, R. D. & A. LATHROP, 2004b. *Micrurus ibiboboca* (Serpentes, Elapidae) is not a Guiana Shield species. **Phyllomedusa** 3(2): 141-144.
- MACCULLOCH, R., R. REYNOLDS, M. T. RODRIGUES & A. MIJARES, 2004. *Pristimantis marmoratus*. In: IUCN 2009. **IUCN Red List of Threatened Species. Version 2009.1**. Disponible in: <[www.iucnredlist.org](http://www.iucnredlist.org)>. Accessed on: 23 August 2009.
- MACCULLOCH, R. D. & A. LATHROP, 2005. Hylid frogs from Mount Ayanganna, Guyana: new species, redescrptions, and distributional records. **Phyllomedusa** 4: 17-37.
- MACCULLOCH, R. D., A. LATHROP & S. Z. KAHN, 2006. Exceptional diversity of *Stefania* (Anura, Cryptobatrachidae) II: six species from Mount Wokomung, Guyana. **Phyllomedusa** 5(1): 31-41.
- MACCULLOCH, R. D., A. LATHROP, R. P. REYNOLDS, J. CELSA SEÑARIS & G. SCHNEIDER, 2007. Herpetofauna of Mount Roraima, Guiana Shield Region, Northeastern South America. **Herpetological Review** 38(1): 24-30.
- MACCULLOCH, R. D., A. LATHROP, P. J. R. KOK, L. R. MINTER, S. Z. KHAN & C. L. BARRIO-AMORÓS, 2008a. A new species of *Adelophryne* (Anura: Eleutherodactylidae) from Guyana, with additional description of *A. guttuosa*. **Zootaxa** 1884: 36-50.
- MACCULLOCH, R. D., A. LATHROP, L. R. MINTER & S. Z. KAHN, 2008b. *Otophryne* (Anura: Microhylidae) from the highlands of Guyana: redescrptions, vocalizations, tadpoles and new distributions. **Papéis Avulsos de Zoologia** 48(22): 247-261.
- MACCULLOCH, R. D. & A. LATHROP, 2009. Herpetofauna of Mount Ayanganna, Guyana. **ROM Contributions in Science** 4: 1-35.
- MARTINS, M. & M. E. OLIVEIRA, 1993. The snakes of the genus *Atractus* Wagler (Reptilia: Squamata) from the Manaus region, central Amazonia, Brazil. **Zoologische Mededelingen Leiden** 67(2): 21-40.
- MARTINS, M. & M. E. OLIVEIRA, 1998. Natural history of snakes in forests of the Manaus region, Central Amazonia, Brazil. **Herpetological Natural History** 6(2): 78-150.
- MASSEMIN, D., D. BORDAGE & K. KUNZ, 2003. Entdeckung einer für Französch-Guyana neuen *Pipa*-Art. **DRACO** 3: 96.
- MASSEMIN, D., D. BORDAGE & K. KUNZ, 2007. Report on the occurrence of *Pipa snethlageae* (Anura: Pipidae) in French Guiana, with notes on its natural history. **Salamandra** 43: 139-147.
- MCDIARMID, R. W. & M. A. DONNELLY, 2005. The Herpetofauna of the Guayana Highlands: Amphibians and Reptiles of the Lost World. In: M. A. DONNELLY, B. I. CROTHER, C. GUYER, M. H. WAKE & M. E. WHITE (Eds.): **Ecology and evolution in the tropics. A herpetological perspective**: 461-560. University of Chicago Press, Chicago.
- MEDEM, F., 1983. **Los Crocodylia de Sur America**. Volumen II: 1-270. Universidad Nacional de Colombia & Colciencias, Bogotá.
- MORALES, V. R., 2002 [2000]. Sistemática y biogeografía del grupo *trilineatus* (Amphibia, Anura, Dendrobatidae, *Colostethus*), com descripción de once nuevas especies. **Publicaciones de la Asociacion de Amigos de Donaña** 13: 1-59.
- MYERS, C. W. & A. S. RAND, 1969. Checklist of amphibians and reptiles of Barro Colorado Island, with comments on faunal change and sampling. **Smithsonian Contributions to Zoology** 10: 1-11.



- MYERS, C. W. & J. E. CADLE, 1994. A new genus for South American snakes related to *Rhadinea obtusa* Cope (Colubridae) and resurrection of *Taeniophallus* Cope for the "*Rhadinea*" *brevirostris* group. **American Museum Novitates** 3102: 1-33.
- NARVAES, P. & M. T. RODRIGUES, 2009. Taxonomic revision of *Rhinella granulosa* species group (Amphibia, Anura, Bufonidae), with a description of a new species. **Arquivos de Zoologia** 40(1): 1-73.
- NOONAN, B. P. & P. GAUCHER, 2005. Phylogeography and demography of Guianan harlequin frogs (*Atelopus*): diversification within a refuge. **Molecular Ecology** 14: 3017-3031.
- NOONAN, B. P. & P. GAUCHER, 2006. Refugial isolation and secondary contact in the dyeing poison frog *Dendrobates tinctorius*. **Molecular Ecology** 15: 4425-4435.
- NOONAN, B. P. & K. P. WRAY, 2006. Neotropical diversification: the effects of a complex history within the poison frog genus *Dendrobates*. **Journal of Biogeography** 33: 1007-1020.
- NUSSBAUM, R. A. & M. S. HOOGMOED, 1979. Surinam Caecilians, with notes on *Rhinatrema bivittatum* and the description of a new species of *Microcaecilia* (Amphibia, Gymnophiona). **Zoologische Mededelingen Leiden** 54(14): 217-235.
- OREJAS-MIRANDA, B., 1967. El genero "*Leptotyphlops*" en el región Amazónica. **Atas do simpósio sobre a Biota Amazónica** 5 (Zoologia): 421-442.
- PARKER, H. W., 1940. Undescribed anatomical structures and new species of reptiles and amphibians. **The Annals and Magazine of Natural History** 11: 257-274.
- PEEL, M. C., B. L. FINLAYSON & T. A. MCMAHON, 2007. Updated world map of the Köppen-Geiger climate classification. **Hydrology and Earth System Sciences** 1: 1633-1644.
- PELOSO, P. L. V. & T. C. S. AVILA-PIRES. Morphological variation in *Ptychoglossus brevifrontalis* Boulenger, 1912 and the status of *Ptychoglossus nicefori* (Loveridge, 1929) (Squamata, Gymnophthalmidae). **Herpetologica** (in press).
- PELOSO, P. L. V. & M. J. STURARO, 2008. A new species of narrow-mouthed frog of the genus *Chiasmocleis* Méhely 1904 (Anura, Microhylidae) from the Amazonian rainforest of Brazil. **Zootaxa** 1947: 39-52.
- PINTO, M. G. M. & W. E. QUATMAN, 2005. Geographic Distribution: *Ptychoglossus brevifrontalis* (Boulenger's Large-scaled Lizard). **Herpetological Review** 36: 202.
- PRITCHARD, P. C. H. & P. TREBBAU, 1984. **The turtles of Venezuela**: 1-399. Society for the Study of Amphibians and Reptiles, Ithaca.
- RIBEIRO-JUNIOR, M. A., T. A. GARDNER & T. C. S. AVILA-PIRES, 2008. Evaluating the effectiveness of herpetofaunal sampling techniques across a gradient of habitat change in a tropical forest landscape. **Journal of Herpetology** 42(4): 733-749.
- RODRIGUES, M. T., R. REYNOLDS & C. GASCON, 2004. *Chiasmocleis shudikarensis*. In: IUCN 2009. **IUCN Red List of Threatened Species. Version 2009.1**. Disponible in: <www.iucnredlist.org>. Accessed on: 23 August 2009.
- RON, S., 2005. Predicting the distribution of the amphibian pathogen *Batrachochytrium dendrobatidis* in the New World. **Biotropica** 37(2): 209-221.
- ROZE, J., 1996. **Coral snakes of the Americas**: i-xii, 1-328. Krieger, Malabar.
- RUEDA-ALMONACID, J. V., J. L. CARR, R. A. MITTERMEIER, J. V. RODRÍGUEZ-MAHECHA, R. B. MAST, R. C. VOGT, A. G. J. RHODIN, J. OSSA-VELÁSQUEZ, J. N. RUEDA & C. G. MITTERMEIER, 2007. **Las Tortugas y los cocodrilianos de los países andinos del trópico**: 1-537. Serie Guías Tropicales de Campo, Conservación Internacional, Bogotá.
- SCOTT, N. J., 1994. Complete species inventories. In: W. R. HEYER, M. A. DONNELLY, R. W. MCDIARMID, L.-A. C. HAYEK & M. S. FOSTER (Eds.): **Measuring and monitoring biological diversity. Standard methods for amphibians**: 84-92. Smithsonian Institution Press, Washington.
- SEBA, A., 1734. *Locupletissimi rerum naturalium thesauri accurata descriptio, et iconibus artificiosissimis expressio, per universam physiceps historiam*. Vol. I: 1-212. J. Wetstenius, G. Smith, J. Waesbergius, Amsterdam.
- SEÑARIS, C. J. & A. ACOSTA-GALVIS, 2004. *Otophryne pyburni*. In: IUCN 2009. **IUCN Red List of Threatened Species. Version 2009.2**. Disponible in: <www.iucnredlist.org>. Accessed on: 17 February 2010.
- SEÑARIS, J. C. & J. AYARZAGÜENA, 2005. **Revisión taxonómica de la Familia Centrolenidae (Amphibia; Anura) de Venezuela**: i-vii, 1-337. Publicaciones del Comité Español del Programa Hombre y Biosfera – Red IberoMaB de la UNESCO, Sevilla.
- SEÑARIS, J. C. & R. MACCULLOCH, 2005. Amphibians. In: T. HOLLOWELL & R. P. REYNOLDS (Eds.): Checklist of the terrestrial vertebrates of the Guiana Shield. **Bulletin of the Biological Society of Washington** 13: 9-23.
- SEÑARIS, J. C., C. A. LASSO, G. RIVAS, M. KALAMANDEEN & E. MANAWANARU, 2008. Amphibians and reptiles of the Acarai Mountains, and Sipu, Kamo and Essequibo rivers in the Konashen COCA, southern Guyana. In: L. A. ALONSO, J. N. MCCULLOUGH, P. NASKRECKI, E. ALEXANDER & W. E. WRIGHT (Eds.): **A rapid biological assessment of the Konashen Community Owned Conservation Area, Southern Guyana**: 55-62. Conservation International (RAP Bulletin of Biological Assessment, 51), Arlington.
- SILVA, J. M. C., A. B. RYLANDS & G. A. B. FONSECA, 2005. The fate of the Amazonian areas of endemism. **Conservation Biology** 19(3): 689-694.
- SILVERSTONE, P. A., 1976. A revision of the poison-arrow frogs of the genus *Phyllobates* Bibron in Sagra (Family Dendrobatidae). **Natural History Museum of Los Angeles County Science Bulletin** 27: 1-53.

- STARACE, F., 1998. **Guide des serpents et amphibènes de Guyane**: 1-449. Ibis Rouge Editions, Guyane.
- STOKSTAD, E., 2008. A second chance for rainforest biodiversity. **Science** 329: 1436-1438.
- SUDAM, 1984. **Atlas climatológico da Amazônia Brasileira**: 1-125. SUDAM (Projeto de hidrologia e climatologia da Amazônia, 19), Belém.
- TRUEB, L. & D. C. CANNATELLA, 1986. Systematics, morphology and phylogeny of genus *Pipa* (Anura: Pipidae). **Herpetologica** 42(4): 412-449.
- TWOMEY, E. & J. L. BROWN, 2008. A partial revision of the *Ameerega hahneli* complex (Anura: Dendrobatidae) and a new cryptic species from the East-Andean versant of Central Peru. **Zootaxa** 1757: 49-65.
- VANZOLINI, P. E. & C. M. CARVALHO, 1991. Two sibling and sympatric species of *Gymnophthalmus* in Roraima, Brasil (Sauria: Teiidae). **Papéis Avulsos de Zoologia** 37(12): 173-226.
- VIDAL, N., M. DEWYNTER & D. GOWER, 2010. Dissecting the major American snake radiation: A molecular phylogeny of the Dipsadidae Bonaparte (Serpentes, Caenophidia). **Comptes Rendus Biologies** 333(2010): 48-55.
- VITT, L. J., W. E. MAGNUSSON, T. C. S. AVILA-PIRES & A. P. LIMA, 2008. **Guia de lagartos da Reserva Adolpho Ducke, Amazônia Central**: 1-175. INPA, Manaus.
- VOGT, R. C., 1994. Reproduction of the cabeçudo, *Peltocephalus dumerilianus*, in the Biological Reserve of Rio Trombetas, Brazil. **Chelonian Conservation and Biology** 1: 159-162.
- VOGT, R. C., 2008. **Amazon turtles**: 1-104. Grafica Biblio, Lima.
- WAGLER, J., 1824. *Serpentum brasiliensium species novae ou Histoire Naturelle des espèces nouvelles de serpents, recueillis et observées pendant le voyage dans l'intérieur du Brésil dans les années 1817, 1818, 1819, 1820 executé par ordre de Sa Majesté le Roi de Bavière, publiée par Jean de Spix, écrite d'après les notes du voyageur*: 1-75. Typis franc. Seraph. Hübschmanni, Munich.
- WOLLENBERG, K. C., M. VEITH, B. P. NOONAN & S. LÖTTERS, 2006. Polymorphism versus species richness – systematic of large *Dendrobates* from the eastern Guiana Shield (Amphibia: Dendrobatidae). **Copeia** 2006(4): 623-629.
- WOLLENBERG, K. C., S. LÖTTERS, C. MORA-FERRER & M. VEITH, 2008. Disentangling composite colour patterns in a poison frog species. **Biological Journal of the Linnean Society** 93: 433-444.
- YANEK, K., W. R. HEYER & R. O. DE SÁ, 2006. Genetic resolution of the enigmatic Lesser Antillean distribution of the frog *Leptodactylus validus* (Anura, Leptodactylidae). **South American Journal of Herpetology** 1(3): 192-201.
- YÁNEZ-MUÑOZ, M., P. PÉREZ-PEÑA & D. CISNEROS-HEREDIA, 2009. New country records of *Hyalinobatrachium iaspidiense* (Amphibia, Anura, Centrolenidae) from the Amazonian lowlands of Ecuador and Peru. **Herpetology Notes** 2: 49-52.
- ZAHER, H., F. G. GRAZIOTTIN, J. E. CADLE, R. W. MURPHY, J. C. MOURA LEITE & S. L. BONATTO, 2009. Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines: a revised classification and descriptions of new taxa. **Papéis Avulsos de Zoologia** 49(11): 115-153.

Recebido: 07/09/2009

Aprovado: 16/03/2010



**APPENDIX 1.** Species of amphibians and reptiles found in the seven studied sites plus Estanifera (Rio Tinto basecamp, where only occasional collections were made) in northern Pará, Brasil, with data on reproduction, diet and habitat based on personal observations and literature. Species indicated with an asterisk are new for the herpetofauna of Brazil, those indicated with a '+' are new for the State of Pará.

(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
Amphibia														
Anura														
<b>Allophrynidae</b>														
<i>Allophryne ruthveni</i> Gauge, 1926					x		x				eggs on leaves	rain forest	arboreal	insects
<b>Bufo</b>														
<i>Ateolopus hoogmoedi</i> Lescure, 1974				x					Guiana south to Tucuruí, Carajás, Santarém		eggs in creeks	rain forest near creeks	terrestrial	insects
<i>Bufo guttatus</i> Schneider, 1799	x	x	x	x	x		x	x			eggs in pools	rain forest	terrestrial	insects
* <i>Bufo</i> sp. n.	x								x		eggs in pools	rain forest	terrestrial	insects
<i>Bufo margaritifera</i> Laurenti, 1768		x		x	x	O	x	x			eggs in pools	rain forest	terrestrial	insects
<i>Bufo marinus</i> Linnaeus, 1758		x		x	x		x	x			eggs in pools	generalist	terrestrial	insects, small vertebrates
<i>Dendrophryniscus bokermanni</i> Izecksohn, 1994	x										terrestrial eggs, edge of pools	rain forest	terrestrial	insects






(Continued)

Diet	insects	insects	insects	insects	insects	insects	insects	insects	insects
Microhabitat	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial
Habitat	rain forest	rain forest edge of openings	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest
Reproductive mode	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles in nest leaves until metamorphosis	terrestrial eggs, one or two tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools	terrestrial eggs, tadpoles on back adult transported to pools
CITES			x	x	x	x	x	x	x
Endemic of Guiana Shield	x	x	x	x	x	x	x	x	x
FLOTA Paru	x								A
REBIO Maicuru	x						x		
Grão-Pará (Estanifera)									
Grão-Pará South									x
Grão-Pará Centre									x
Grão-Pará North	x		x	x	x	x	x	x	x
FLOTA Trombetas	x						x	x	x
FLOTA Faro		x							
Species name	<i>Anomaloglossus baobatrachus</i> (Boistel & Massary, 1999)	<i>Anomaloglossus stephni</i> (Martins, 1989)	<i>Dendrobates tinctorius</i> (Cuvier, 1797)	* <i>Epipedobates</i> cf. <i>guayanensis</i> Heatwole, Solano & Heatwole, 1965	<i>Epipedobates hahnelti</i> (Boulenger, 1884)	<i>Epipedobates trivittatus</i> (Spix, 1824)			



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<b>Hylidae</b>														
<i>Hyla boans</i> (Linnaeus, 1758)		A	×	×			×	×			nests at edge of rivers, creeks and pools	rain forest, gallery forest	arboreal	insects
<i>Hyla calcarata</i> Troschel, 1848			×		×		×				eggs in water	creeks, flooded areas, forest and edge	arboreal	insects
<i>Hyla cinerascens</i> Spix, 1824			×								eggs in lentic water	rain forest, savanna forest, close to creeks, flooded areas	arboreal	insects
+ <i>Hyla dentei</i> Bokermann, 1967							×		×		?	rain forest	arboreal	insects
<i>Hyla fasciata</i> (Günther, 1858)					×		×	×			eggs in water	rain forest and edge	arboreal	insects
* <i>Hyla gaucheri</i> Lescure & Marty, 2000					×				×		?	pools in rain forest and open areas	arboreal	insects
<i>Hyla geographica</i> Spix, 1824				×	×		×				eggs in lentic or lotic water	savanna, open areas, rain forest	arboreal	insects
<i>Hyla leucophyllata</i> (Beireis, 1783)					×		×				eggs on leaves above pool	open areas, savanna, rain forest and edge	arboreal	insects




(Continued)

Diet	insects	insects	insects	insects	insects	insects	insects	insects	insects
Microhabitat	arboreal	arboreal	arboreal	arboreal	arboreal	arboreal	arboreal	arboreal	arboreal
Habitat	rain forest, savanna, open areas	generalist (open areas)	generalist (open areas)	savanna, open areas, rain forest	open areas / antropic, savanna, swamps	flooded riverine forest (várzea)	savanna, swamps, rain forest	rain forest	rain forest
Reproductive mode	eggs in water	eggs on leaves above pool	eggs above or in pools (?)	eggs on leaves above lentic water	eggs in water	eggs in nest at edge of water	eggs in water	eggs in Bromeliaceae with parental care	
CITES									
Endemic of Guiana Shield									
FLOTA Paru			x		x		x		
REBIO Maicuru							x		
Grão-Pará (Estanífera)									
Grão-Pará South			x	x					x
Grão-Pará Centre	x	x	x				x		
Grão-Pará North									x
FLOTA Trombetas		x	A						x
FLOTA Faro						x	x		
Species name	<i>Hyla minuscula</i> Rivero, 1971	<i>Hyla minuta</i> Peters, 1872	<i>Hyla multifasciata</i> Gunther, 1859	<i>Hyla</i> cf. <i>nana</i> Boulenger, 1889	<i>Hyla punctata</i> (Schneider, 1799)	<i>Hyla warrini</i> Parker, 1936	<i>Osteocephalus lepriouri</i> (Duméril & Bribron, 1841)	<i>Osteocephalus oophagus</i> Jungfer & Schiesari, 1995	



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Osteocephalus taurinus</i> Steindachner, 1862	x	x	x	x	x		A			eggs in temporary pools	rain forest	arboreal	insects
<i>Phyllomedusa bicolor</i> (Boddaert, 1772)						x				nest between leaves above water	savanna, rain forest, várzea	arboreal	insects
<i>Phyllomedusa hypochondrialis</i> (Daudin, 1803)		A		x		x				nest between leaves above water	generalist (open areas)	arboreal	insects
<i>Phyllomedusa tomopterna</i> (Cope, 1868)		A		x	A					nest between leaves above water	rain forest	arboreal	insects
<i>Phyllomedusa vaillanti</i> Boulenger, 1882										nest between leaves above water	rain forest	arboreal	insects
* <i>Phrynohyas hadroceps</i> (Duellman & Hoogmoed, 1992)						A		x		eggs in cavities in trees, above the ground	rain forest	arboreal	insects
<i>Phrynohyas</i> sp.							A					arboreal	
<i>Scinax boesemani</i> Goin, 1966										eggs in lentic water	savanna, open areas, rain forest in open areas, várzea	arboreal	insects
+ <i>Scinax garbei</i> (Miranda-Ribeiro, 1926)						x				eggs in lentic water	rain forest	arboreal	insects



(Continued)

Diet	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects, small vertebrates	insects
Microhabitat	arboreal	arboreal	arboreal	arboreal	arboreal		terrestrial	terrestrial	terrestrial	semifossorial	arboreal		
Habitat	open areas, savanna	rain forest	perianthropic, savanna	rain forest	rain forest		rain forest	rain forest	savanna	rain forest	rain forest		
Reproductive mode	eggs in lentic water	eggs in lentic water	eggs in lentic water	eggs in lentic water	eggs in lentic water		nest in ground	nest in ground	nest in ground	eggs in lentic water	terrestrial eggs, direct development		
CITES													
Endemic of Guiana Shield		x						x			x		
FLOTA Paru					x		x						
REBIO Maicuru				x			x				x		
Grão-Pará (Estanifera)													
Grão-Pará South		x	x				x			x			
Grão-Pará Centre	x						x		x				
Grão-Pará North							x	x					
FLOTA Trombetas							x				x		
FLOTA Faro							x						
Species name	<i>Scinax nebulosus</i> (Spix, 1824)	+ <i>Scinax proboscideus</i> (Brongersma, 1933)	<i>Scinax ruber</i> (Laurenti, 1768)	* <i>Scinax</i> sp. nov.	<i>Scinax</i> gr. <i>ruber</i> (Laurenti, 1768)	<b>Leptodactylidae</b>	<i>Adenomera andreae</i> (Müller, 1923)	* <i>Adenomera heyeri</i> Boistel, Massary & Angulo, 2006	<i>Adenomera hylaedactylia</i> (Cope, 1868)	<i>Ceratophrys cornuta</i> (Linnaeus, 1758)	+ <i>Eleutherodactylus chiastonotus</i> Lynch & Hoogmoed, 1977		



(Continued)

Diet	insects	insects	insects	insects	insects, small vertebrates	insects, small vertebrates	insects	insects
Microhabitat	arboreal	arboreal	arboreal	arboreal	terrestrial	terrestrial	terrestrial	terrestrial
Habitat	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	savanna
Reproductive mode	terrestrial eggs, direct development	terrestrial eggs, direct development	terrestrial eggs, direct development	terrestrial eggs, direct development	foamnest	terrestrial foamnest	terrestrial foamnest	terrestrial foamnest in tunnel
CITES								
Endemic of Guiana Shield		x	x					nearly
FLOTA Paru					x	x		
REBIO Maicuru	?	x			x	x		
Grão-Pará (Estanifera)								
Grão-Pará South				x	x	x		
Grão-Pará Centre				x		x		x
Grão-Pará North	x	x	x	x		x	x	
FLOTA Trombetas				x	x	x		
FLOTA Faro				x		x		
Species name	+ <i>Eleutherodactylus fenestratus</i> (Steindachner, 1864)	+ <i>Eleutherodactylus inguinalis</i> Parker, 1940	+ <i>Eleutherodactylus marmoratus</i> (Boulenger, 1900)	<i>Eleutherodactylus zeuctotylus</i> Lynch & Hoogmoed, 1977	+ <i>Leptodactylus bolivianus</i> Boulenger, 1898	<i>Leptodactylus knudseni</i> Heyer, 1972	<i>Leptodactylus leptodactyloides</i> (Andersson, 1945)	<i>Leptodactylus longirostris</i> Boulenger, 1882



(Continued)

Diet	insects	insects	insects; small vertebrates	insects	insects	insects	insects	insects	insects
Microhabitat	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial	terrestrial
Habitat	open rockslates in savanna and rain forest	rain forest	rain forest	generalist (open areas)	rain forest	rain forest	rain forest	rain forest	rain forest
Reproductive mode	terrestrial foamnest under stones	terrestrial foamnest	terrestrial foamnest	foamnest at edge water	foamnest at edge water	terrestrial foamnest	foamnest in <i>Atta</i> ants nest	foamnest floating on water	
CITES									
Endemic of Guiana Shield	x								Guiana + Belém
FLOTA Paru		x		x			x		
REBIO Maicuru				x			x	x	
Grão-Pará (Estanífera)									x
Grão-Pará South		x	x	x	x		x	x	
Grão-Pará Centre	x	x	x		x		x	x	
Grão-Pará North			x				x		
FLOTA Trombetas	x	x	x	x		x	x	x	
FLOTA Faro		x	x	x	x		x		
Species name	<i>Leptodactylus myersi</i> Heyer, 1995	<i>Leptodactylus mystaceus</i> (Spix, 1824)	<i>Leptodactylus pentadactylus</i> (Laurenti, 1768)	<i>Leptodactylus petersii</i> (Steindachner, 1864)	<i>Leptodactylus rhodomystax</i> Boulenger, 1884	<i>Leptodactylus stenodema</i> Jimenez de la Espada, 1875	<i>Lithodytes lineatus</i> (Schneider, 1799)	<i>Physalaemus ephippifer</i> (Steindachner, 1864)	



(Continued)

Diet	Microhabitat	Habitat	Reproductive mode	CITES	Endemic of Guiana Shield	FLOTA Paru	REBIO Maicuru	Grão-Pará (Estanifera)	Grão-Pará South	Grão-Pará Centre	Grão-Pará North	FLOTA Trombetas	FLOTA Faro	Species name
														<b>Microhylidae</b>
insects	fossorial	rain forest	terrestrial eggs; tadpoles in lentic water		x				x				x	+ <i>Chiasmocleis hudsoni</i> Parker, 1940
insects	fossorial	rain forest	?			x	x			x				* <i>Chiasmocleis</i> sp. nov.
insects	fossorial	rain forest	eggs in lentic water			x			x					<i>Hemiphrone boliviana</i> (Parker, 1927)
insects	fossorial	rain forest, near creeks	eggs in lotic water; tadpoles in substrate		x						x			<i>Otophryne pyburni</i> Campbell & Clarke, 1998
insects	fossorial	rain forest, under layer of roots	Eggs subterraneous, direct development		Guiana + eastern Colombia						x	x	x	+ <i>Synapturanus mirandaireiroi</i> Nelson & Lescure, 1975
														<b>Pipidae</b>
aquatic invertebrates and small fish	aquatic	in pools, creeks, rivers, swamps, periantropic	eggs in back mother, direct development			x	x							<i>Pipa pipa</i> (Linnaeus, 1758)
aquatic invertebrates	aquatic	in pools and creeks in rain forest	eggs in back mother; tadpoles direct development										x	<i>Pipa snethlageae</i> Müller, 1914



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanífera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<b>Ranidae</b>						x					eggs in water	rain forest near creeks, savanna forest near pools	terrestrial	insects
+ <i>Rana palmipes</i> Spix, 1824														
<b>GYMNOPHIONA</b>														
<b>Caeciliidae</b>														
<i>Caecilia tentaculata</i> (Linnaeus, 1758)		x		x							juveniles aquatic	rain forest	fossorial	?
* <i>Microcaecilia unicolor</i> (Duméril, 1863)		x							x		?	rain forest	fossorial	?
<b>Rhinatrematidae</b>														
* <i>Rhinatrema</i> sp. nov.									x		?	rain forest	fossorial	?
<b>REPTILIA</b>														
<b>SQUAMATA</b>														
<b>LIZARDS</b>														
<b>Gekkonidae</b>														
<i>Coleodactylus amazonicus</i> (Andersson, 1918)		x		x				x			eggs (one at a time)	rain forest, leaf litter sunny areas	leaf litter	insects
<i>Gonatodes annularis</i> Boulenger, 1887		x							x		eggs (two at a time)	rain forest, base of trees, rocks	arboreal	insects



(Continued)

Diet	insects	insects	insects	insects		insects	insects	insects	insects
Microhabitat	arboreal	leaf litter	leaf litter	arboreal, nocturnal		leaf litter	leaf litter	leaf litter	leaf litter
Habitat	rainforest, on small trees and branches	rain forest, leaf litter near moist areas	rainforest, leaf litter	rain forest on tree trunks		rain forest	rain forest	rain forest	rain forest
Reproductive mode	eggs (two at a time)	eggs (one at a time)	eggs (one at a time)	eggs (two at a time)		eggs	eggs	eggs	eggs
CITES									
Endemic of Guiana Shield						x	Guiana, Belém, Xingu,		
FLOTA Paru	x					x			x
REBIO Maicuru	x	x	x			x	x		x
Grão-Pará (Estanifera)									
Grão-Pará South	x			x					
Grão-Pará Centre	○								
Grão-Pará North	x		x	x			x	x	
FLOTA Trombetas	x	x					x		
FLOTA Faro	x	x	x						x
Species name	<i>Gonatodes humeralis</i> (Guichenot, 1855)	<i>Lepidoblepharis heyerorum</i> Vanzolini, 1978	<i>Pseudogonatodes guianensis</i> Parker, 1935	<i>Thecadactylus rapicauda</i> (Houttuyn, 1782)	<b>Gymnophthalmidae</b>	<i>Alopoglossus angulatus</i> (Linnaeus, 1758)	+ <i>Amapasaurus tetradactylus</i> Cunha, 1970	<i>Artrosaura kockii</i> (Lidth de Jeude, 1904)	<i>Artrosaura reticulata</i> (O'Shaughnessy, 1881)



(Continued)

Diet	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects	insects
Microhabitat	leaf litter	leaf litter	leaf litter	leaf litter	terrestrial, under rocks	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter	leaf litter
Habitat	rain forest	rain forest	rain forest	rain forest	savanna	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest	rain forest, creeks				
Reproductive mode	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs
CITES																			
Endemic of Guiana Shield		x			x				Guiana, Belém					Guiana, Belém, Xingu				x	
FLOTA Paru	x		x	x					x		x								
REBIO Maicuru	x		x						x										
Grão-Pará (Estanífera)																			
Grão-Pará South	x								x	x									
Grão-Pará Centre	x			x	x				x	x	x				x				
Grão-Pará North	x			x					x	x	x				x				x
FLOTA Trombetas	x									x									
FLOTA Faro	x	x								x	x								
Species name	<i>Bachia flavescens</i> (Bonnaterre, 1789)	+ <i>Bachia panoplia</i> Thomas, 1969	<i>Cercosaura argulus</i> Peters, 1863	<i>Cercosaura ocellata</i> Wagler, 1830	<i>Gymnophthalmus</i> cf. <i>underwoodi</i> Grant, 1958	<i>Iphisa elegans</i> Gray, 1851	<i>Leposoma guianense</i> Ruibal, 1952	<i>Leposoma</i> <i>percarinatum</i> Muller, 1923)	<i>Leposoma</i> sp. nov.	<i>Neusticurus</i> <i>bicarinatus</i> (Linnaeus, 1758)	<i>Neusticurus rudis</i> Boulenger, 1900								



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanífera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Neusticurus</i> indet.								○			eggs	rain forest, creeks	semi-aquatic	insects
+ <i>PtychoGLOSSUS brevifrontalis</i> Boulenger, 1912	×		×								eggs	rain forest	leaf litter	insects
<i>Tretioscincus agilis</i> (Ruthven, 1916)	×	×	×				×	○	Guiana, Belém, Xingu		eggs	rain forest, lianas, rotten trees	terrestrial and, arboreal	insects
<b>Iguanidae</b>														
<i>Iguana iguana</i> (Linnaeus, 1758)							○			×	eggs	forest and open areas, on ground or on vegetation, usually near water	arboreal	leaves, flowers, fruits
<b>Polychrotidae</b>														
<i>Anolis auratus</i> Daudin, 1802				×							eggs	savanna, herbaceous vegetation	arboreal	insects
<i>Anolis chrysolepis</i> Duméril & Bibron, 1837	×	×	×				×		×		eggs	rain forest, leaf litter	arboreal	insects
<i>Anolis fuscoauratus</i> d'Orbigny, 1837	×	×	×				×				eggs	rain forest and edge, on vegetation	arboreal	insects
<i>Anolis ortonii</i> Cope, 1868							○				eggs	rain forest and edge on vegetation	arboreal	insects




(Continued)

Diet	insects			insects mainly ants	insects mainly ants	insects			insects
Microhabitat	arboreal			arboreal	arboreal	arboreal			terrestrial and arboreal
Habitat	rain forest on tree trunks			rain forest on large tree trunks	rain forest, secondary forest on small trees	rain forest along margins creeks and várzea along rivers			open areas in rain forest
Reproductive mode	eggs			eggs	eggs	eggs			viviparous
CITES									
Endemic of Guiana Shield									
FLOTA Paru				O	x	x			x
REBIO Maicuru				P	x	x			x
Grão-Pará (Estanífera)									
Grão-Pará South	x				x				
Grão-Pará Centre					P	x			P
Grão-Pará North				x	x	x			x
FLOTA Trombetas					x	x			x
FLOTA Faro					x	x			
Species name	<i>Anolis punctatus</i> Daudin, 1802			<i>Plica plica</i> (Linnaeus, 1758)	<i>Plica umbra</i> (Linnaeus, 1758)	<i>Uranoscodon superciliosus</i> (Linnaeus, 1758)			<i>Mabuya nigropunctata</i> (Spix, 1825)
	<b>Tropiduridae</b>								<b>Teiidae</b>



(Continued)

Diet	insects, small vertebrates	insects, small vertebrates	insects				invertebrates and small vertebrates	invertebrates	invertebrates
Microhabitat	terrestrial	terrestrial and on low vegetation	terrestrial and on low vegetation				fossorial	fossorial	fossorial
Habitat	generalist, open areas, savanna, perianthropic, in rain forest in open areas	rain forest, secondary forest, in open areas	savanna, floating mats of vegetation				rain forest and open vegetation	rain forest	rain forest
Reproductive mode	eggs	eggs	eggs				eggs	eggs	eggs
CITES									
Endemic of Guiana Shield			Guiana + NW South America					x	x
FLOTA Paru	x	x							
REBIO Maicuru	x	x							
Grão-Pará (Estanífera)	P								
Grão-Pará South	x	O					x	x	
Grão-Pará Centre	x	x	x						
Grão-Pará North	O	O							
FLOTA Trombetas	P	x	O						
FLOTA Faro	x	x							x
Species name	<i>Ameiva ameiva</i> (Linnaeus, 1758)	<i>Kentropyx calcarata</i> Spix, 1825	<i>Kentropyx striata</i> (Daudin, 1802)	AMPHISBAENIA	AMPHISBAENIDAE	<i>Amphisbaena fuliginosa</i> Linnaeus, 1758	<i>Amphisbaena vanzolinii</i> Gans, 1963	* <i>Mesobaena rhachicephala</i> Hoogmoed, Pinto, Rocha & Pereira, 2009	



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
SERPENTES														
<b>Anomalepididae</b>														
<i>Typhlops squamosus</i> (Schlegel, 1839)				○					Guiana, Trinidad, Belém		eggs	rain forest	fossorial	ants, termites
<b>Leptotyphlopidae</b>														
<i>Leptotyphlops albifrons</i> (Wagler, 1824)				×				×	×		eggs	rain forest	fossorial	termites, ants
+ <i>Leptotyphlops cupirensis</i> Bailey & Carvalho, 1946					×						eggs	rain forest	fossorial	ants, termites
<b>Typhlopidae</b>														
<i>Typhlops brongersmianus</i> Vanzolini, 1976	×										eggs	rain forest	fossorial	insects
<b>Boidae</b>														
<i>Boa constrictor</i> Linnaeus, 1758	○									×	viviparous	rain forest, savanna, periantropic	terrestrial	vertebrates
<i>Corallus caninus</i> (Linnaeus, 1758)									×	×	viviparous	rain forest	arboreal	vertebrates
<i>Corallus hortulanus</i> (Linnaeus, 1758)	×						×			×	viviparous	along rivers and creeks in rain forest and savanna	arboreal	small vertebrates



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Eumeces murinus</i> (Linnaeus, 1758)				x						x	viviparous	in rivers and creeks in rain forest, savanna, swamps, periantropic	aquatic	vertebrates
<b>Colubridae</b>														
+ <i>Apostolepis nigrolineata</i> (Peters, 1869)			x	x							eggs	rain forest	fossorial	?
<i>Apostolepis quinque-lineata</i> Boulenger, 1896			x		x						eggs	rain forest	fossorial	?
+ <i>Atractus badius</i> (F.Boie, 1827)					x						eggs	rain forest	fossorial	probably earthworms
<i>Atractus torquatus</i> (Duméril, Bibron & Duméril, 1854)			x								eggs	rain forest	fossorial	
<i>Chironius exoletus</i> (Linnaeus, 1758)	x										eggs	rain forest	terrestrial	small vertebrates
<i>Chironius fuscus</i> (Linnaeus, 1758)			x								eggs	rain forest, savanna, periantropic	terrestrial	small vertebrates



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Chironius multiventris</i> <i>cochranae</i> Hoge & Romano, 1969		x									eggs	rain forest	terrestrial	small vertebrates
<i>Chironius scurrulus</i> (Wagler, 1824)			x					x			eggs	rain forest, open areas	terrestrial	amphibians and reptiles
<i>Clelia clelia</i> (Daudin, 1803)		x								x	eggs	rain forest	terrestrial	snakes
<i>Dendrophidion</i> <i>dendrophis</i> (Schlegel, 1837)			x				x				eggs	rain forest	terrestrial	small anurans
<i>Dipsas catesbyi</i> (Sentzen, 1796)		x					x				eggs	rain forest	arboreal	mollusks
<i>Drymoluber dichrous</i> (Peters, 1863)		x						x			eggs	rain forest	terrestrial	lizards
<i>Erythrolamprus</i> <i>aesculapii</i> (Linnaeus, 1766)			x				x	○			eggs	rain forest	terrestrial	snakes
<i>Helicops angulatus</i> (Linnaeus, 1758)		x						x			viviparous	creeks, pools, aquatic generalist	aquatic	fishes, amphibians
<i>Imantodes cenchoa</i> (Linnaeus, 1758)		x									eggs	rain forest	arboreal	lizards, treefrogs
<i>Leptodeira annulata</i> (Linnaeus, 1758)				x			x				eggs	rain forest, savanna, open areas, perianthropic	arboreal	anurans and eggs of anurans



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Leptophis ahaetulla</i> (Linnaeus, 1758)		P									eggs	rain forest, savanna, open areas, periantropic	arboreal	small vertebrates
<i>Liophis reginae</i> (Linnaeus, 1758)			x		x		x				eggs	rain forest, periantropic near water	terrestrial	anurans
<i>Liophis typhlus</i> (Linnaeus, 1758)		x		P				x			eggs	rain forest	terrestrial	anurans
<i>Mastigodyas boddaerti</i> (Sentzen, 1796)		x			x			x			eggs	generalist	terrestrial	small vertebrates
<i>Oxyrhopus petola</i> (Linnaeus, 1758)								x			eggs	rain forest and secondary forest, savanna	terrestrial	small mammals
<i>Pseudoboa coronata</i> Schneider, 1801											eggs	rain forest and secondary forest, periantropic	terrestrial	small mammals reptiles, birds
<i>Pseudoboa newwedii</i> (Duméril, Bibron & Duméril, 1854)											eggs	savanna	terrestrial	reptiles, small mammals
<i>Pseustes poecilonotus</i> Günther, 1858											eggs	rain forest and secondary forest, periantropic	terrestrial	mammals, birds, lizards



(Continued)

Diet	mammals, birds, lizards	birds, mammals	anurans, lizards, mammals	scolopenders	small lizards	small frogs	anurans	anurans and tadpoles
Microhabitat	terrestrial	terrestrial	arboreal	fossorial	terrestrial	terrestrial	terrestrial	terrestrial
Habitat	rain forest and secondary forest, periantropic	rain forest	rain forest	rain forest	rain forest and secondary forest, periantropic	rain forest	rainforest near creeks	rain forest and secondary forest
Reproductive mode	eggs	eggs	eggs	eggs	eggs	eggs	eggs	eggs
CITES								
Endemic of Guiana Shield						x	x	
FLOTA Paru								
REBIO Maicuru	x							x
Grão-Pará (Estanifera)								
Grão-Pará South	x			x				
Grão-Pará Centre	x	x	x					
Grão-Pará North	x				x	x	x	
FLOTA Trombetas								
FLOTA Faro				x	x			
Species name	<i>Pseustes sulphureus</i> (Wagler, 1824)	<i>Rhinobothryum leniginosum</i> (Scopoli, 1785)	<i>Siphophis compressus</i> (Daudin, 1803)	<i>Tantilla melanocephala</i> (Linnaeus, 1758)	<i>Taeniophallus brevirostris</i> (Peters, 1863)	+ <i>Taeniophallus nicagus</i> (Cope, 1863)	+ <i>Thalesius viridis</i> (Werner, 1924)	<i>Xenopholis scalaris</i> (Wucherer, 1861)



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	Grão-Pará (Estanifera)	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Xenoxylis argenteus</i> (Daudin, 1803)			x								eggs	rain forest and secondary forest	arboreal	lizards, anurans, birds
<b>Elapidae</b>														
+ <i>Micrurus averyi</i> Schmidt, 1939			x						x		eggs	rain forest	terrestrial	?
<i>Micrurus hemprichii</i> (Jan, 1858)				x							eggs	rain forest	terrestrial	Onychophora
<i>Micrurus lemniscatus</i> (Linnaeus, 1758)	x						x				eggs	savanna, rain forest	terrestrial	fishes, aquatic snakes
<i>Micrurus paraensis</i> Cunha & Nascimento, 1973		○			x				x		eggs	rain forest	terrestrial	small snakes
<i>Micrurus surinamensis</i> (Cuvier, 1817)	x										eggs	in open areas, rain forest	aquatic	fishes
<b>Viperidae</b>														
<i>Bothriopsis taeniata</i> (Wagler, 1824)		x									viviparous	rain forest	arboreal	small vertebrates
<i>Bothrops atrox</i> (Linnaeus, 1758)		x	x					x			viviparous	rain forest and secondary forest	terrestrial	small vertebrates



(Continued)

Species name	FLOTA Faro	FLOTA Trombetas	Grão-Pará North	Grão-Pará Centre	Grão-Pará South	REBIO Maicuru	FLOTA Paru	Endemic of Guiana Shield	CITES	Reproductive mode	Habitat	Microhabitat	Diet
<i>Lachesis muta</i> (Linnaeus, 1766)						x	x			eggs	rain forest	terrestrial	small vertebrates
<b>CHELONIA<sup>1</sup></b>													
<b>Testudinidae</b>													
<i>Chelonoidis carbonaria</i> (Spix, 1824)	x	P			x				x	eggs	savannas and rain forest	terrestrial	herbivorous, coprophagous
<i>Chelonoidis denticulata</i> (Linnaeus, 1766)			x	P	x	x	P		x	eggs on forest floor	forest, sometimes in savanna	terrestrial	omnivorous
<b>Chelidae</b>													
<i>Mesoclemmys gibba</i> (Schweigger, 1812)										eggs	creeks in rain forest	aquatic	fishes, tadpoles, aquatic invertebrates, vegetable matter
<i>Platemys platycephala</i> (Schneider, 1792)		x								eggs	creeks and pools in rain forest	aquatic	fishes, tadpoles, aquatic invertebrates

<sup>1</sup> While this article was in the proof stage, we received some pictures from Luciano FA. Montag from which it is clear that *Poecocnemis unifilis* and *Chelonoidis denticulata* also were observed (P) in FLOTA Faro. We have not included these species in Appendix 1 or in any of the tables, graphs or computations. The number of species of reptiles observed in FLOTA Faro consequently increases with two and the total number of species of reptiles collected/observed during the Calha Norte expeditions increases with one. The total number of species of reptiles known for northern Pará does not change because both species had been reported from other localities in the area already.





**APPENDIX 2.** Specimens collected during the expeditions to CNP and deposited in the herpetological collection of MPEG (CN = field numbers). ESEC = Estação Ecológica (Ecological Station).

AMPHIBIA. *Anura*. **Allophryinae**. *Allophryne ruthveni* - ESEC Grão-Pará South: CN850. REBIO Maicuru: CN1381, CN1603, CN1604, CN1630, CN1658, CN1671. **Bufo**. *Atelopus hoogmoedi* - ESEC Grão-Pará Centre: CN2088, CN2097-2102, CN2130-2132, CN2153-2166, CN2237-2239, CN2277, CN2297, CN2299, 2370. *Bufo guttatus* - FLOTA Faro: CN55, CN115, CN207, CN234, CN299, CN312. FLOTA Trombetas: CN 410. ESEC Grão-Pará South: CN831. REBIO Maicuru: CN1397, CN1412, CN1429, CN1440, CN1470-1475, CN1592, CN1617, CN1620, CN1636, CN1637, CN1693. FLOTA Paru: CN1842, CN1854, CN1931, CN1939, CN1998-2000, CN2045, CN2046. ESEC Grão-Pará Centre: CN2188, CN2262, CN2365. *Bufo* sp. n.- FLOTA Faro: CN23, CN24, CN26, CN32, CN33, CN38, CN40, CN41, CN44, CN46, CN48, CN50, CN51, CN56, CN58, CN61, CN67, CN70, CN71-75, CN77, CN93, CN99, CN104, CN119, CN125, CN133, CN134, CN136, CN138, CN144-146, CN156, CN169, CN182, CN184, CN187, CN196, CN199, CN226-229, CN236, CN301, CN304. ESEC Grão-Pará North: CN982, CN984, CN995-997, CN1015, CN1016, CN1019, CN1024, CN1035, CN1044, CN1074, CN1081, CN1087, CN1097-1099, CN1115, CN1118, CN1125-1127, CN1141-1143, CN1154, CN1163, CN1167, CN1173-1175, CN1180, CN1181, CN1186, CN1188, CN1191, CN1192, CN1197, CN1210, CN1211-1214, CN1217, CN1219, CN1220, CN1231, CN1257, CN1264, CN1267, CN1270, CN1271, CN1273, CN1276, CN1278, CN1289, CN1290, CN1293, CN1297, CN1303-1305, CN1310, CN1316, CN1320, CN1321, CN1323, CN1324, CN1342, CN1343, CN1353, CN1357. *Bufo margaritifera* - FLOTA Trombetas: CN333, CN339, CN359, CN360, CN418-421, CN426, CN433, CN434, CN474, CN475, CN490-492, CN496, CN515-521, CN539, CN541, CN542, CN559, CN594. ESEC Grão-Pará South: CN617, CN636, CN637, CN640-642, CN645, CN649, CN672, CN690, CN700, CN703-708, CN722, CN729, CN733, CN736, CN737, CN742, CN743, CN765, CN766, CN770, CN781, CN784, CN789, CN794, CN800, CN807, CN809, CN828, CN829, CN849, CN868, CN869, CN870, CN871, CN880, CN882-884, CN899, CN906, CN909, CN921, CN922, CN939, CN942-944, CN965, CN967, CN968; tadpoles CN769, CN838. REBIO Maicuru: CN1396, CN1444, CN1494. FLOTA Paru: CN1703, CN1709, CN1717, CN1724-1735, CN1742, CN1760, CN1796, CN1797, CN1843, CN1876, CN1885, CN1892, CN1893, CN1897, CN1910, CN1933, CN2041, CN2043. ESEC Grão-Pará Centre: CN2055, CN2058, CN2060, CN2061, CN2120-2123, CN2126, CN2142, CN2169, CN2171, CN2182, CN2198, CN2200, CN2201, CN2224, CN2240-2243, CN2248, CN2280, CN2281, CN2301, CN2302, CN2311. *Bufo marinus* - FLOTA Faro: CN47, CN53, CN68, CN92, CN94, CN102, CN103, CN116, CN175, CN246, CN285, CN289, CN290, CN314. FLOTA Trombetas: CN570. ESEC Grão-Pará South: CN620, CN643. REBIO Maicuru: CN1477, CN1607, CN1680. FLOTA Paru: CN1696, CN1700-1702, CN1708, CN1716, CN1720, CN1736, CN1758, CN1761, CN1763, CN1766, CN1779, CN1803, CN1818, CN1827, CN1867, CN1895, CN1937, CN1944, CN1973, CN2030, CN2044. ESEC Grão-Pará Centre: CN2141. *Dendrophryniscus bokermanni* - FLOTA Faro: CN8, CN10, CN20, CN21, CN28, CN29, CN37, CN45, CN65, CN66, CN95, CN98, CN141, CN202, CN203, CN308. *Dendrophryniscus minutus* - FLOTA Faro: CN9, CN11-19, CN43, CN63, CN91, CN96, CN97, CN100, CN105, CN106, CN140, CN186, CN200, CN201. ESEC Grão-Pará North: CN1033, CN1093, CN1112, CN1279, CN1243, CN1279, CN1335, CN1370. REBIO Maicuru: CN1501, CN1529. **Centrolenidae**. *Cochranella* sp.- ESEC Grão-Pará Centre: CN2363. *Hyalinobatrachium iaspidiense*



- ESEC Grão-Pará Centre: CN2362. Girinos indeterminados  
 - ESEC Grão-Pará South: CN684A, CN725A. ESEC Grão-Pará North: CN1137. ESEC Grão-Pará Centre: CN2372.

**Dendrobatidae.** *Allobates femoralis* - FLOTA Trombetas: CN367, CN395, CN480. ESEC Grão-Pará South: CN715. ESEC Grão-Pará North: CN1187, CN1319. REBIO Maicuru: CN1531. *Allobates spumaponens* - ESEC Grão-Pará South: CN628, CN776, CN867, CN935, CN949, CN952. ESEC Grão-Pará North: CN1114, CN1198, CN1237, CN1360, CN1372. REBIO Maicuru: CN1437, CN1560, CN1568, CN1569, CN1624, CN1648. FLOTA Paru: CN1846, 1941. ESEC Grão-Pará Centre: CN2192, CN2193, CN2232, CN2303. *Anomaloglossus baeobatrachus* - FLOTA Trombetas: CN364, CN378, CN449, CN476, CN549. ESEC Grão-Pará North: CN980, CN983, CN985, CN991, CN1004, CN1014, CN1020, CN1022, CN1052-1056, CN1059-1061, CN1066, CN1067-1071, CN1077, CN1078, CN1082-1084, CN1105, CN1106, CN1110, CN1144, CN1155, CN1164, CN1169, CN1195, CN1224, CN1225, CN1229, CN1233, CN1239, CN1244, CN1245, CN1281, CN1285, CN1286, CN1301, CN1312, CN1330, CN1331, CN1355, CN1356, CN1359, CN1371, CN1373. REBIO Maicuru: CN1497-1500, CN1515-1527, CN1530. FLOTA Paru: CN1948-1951, CN1955-1962. *Anomaloglossus stepheni* - FLOTA Faro: CN6, CN204, CN205, CN242, CN251, CN261, CN266, CN267, CN270, CN277, CN278, CN280, CN281, CN287, CN302. *Dendrobates tinctorius* - ESEC Grão-Pará North: CN974, CN975, CN978, CN981, CN986, CN987, CN990, CN1001-1003, CN1005-1010, CN1013, CN1018, CN1027-1029, CN1031, CN1032, CN1034, CN1038, CN1043, CN1046, CN1050, CN1057, CN1058, CN1063, CN1064, CN1085, CN1119-1123. *Epipedobates cf. guayanensis* - FLOTA Trombetas: CN350, CN376, CN390, CN394, CN412, CN425, CN511, CN527, CN533, CN571. *Epipedobates hahneli* - FLOTA Trombetas: CN495. REBIO Maicuru: CN1575. *Epipedobates trivittatus* - FLOTA Trombetas: CN348, CN354, CN361, CN366, CN384, CN403, CN411, CN414, CN415, CN455, CN479, CN488, CN504, CN522, CN546, CN548, CN587, CN596. ESEC Grão-Pará South: CN624, CN795, CN833, CN848. ESEC Grão-Pará Centre: CN2137, CN2190, CN2191, CN2292, CN2359. **Hylidae.** *Hyla boans* - ESEC Grão-Pará North: CN1345. REBIO Maicuru: CN1590. FLOTA Paru: CN1995-1997. ESEC Grão-Pará Centre: CN2081, CN2350. *Hyla calcarata* - ESEC Grão-Pará South: CN685, CN750, CN891-893. REBIO Maicuru: CN1589. *Hyla cinerascens* - ESEC Grão-Pará South: CN684D, CN758 (all tadpoles). *Hyla dentei* - REBIO Maicuru: CN1492, CN1581. *Hyla fasciata* - ESEC Grão-Pará South: CN888, CN889, CN894, CN895. REBIO Maicuru: CN1478, CN1479, CN1582-1588. FLOTA Paru: CN1954. *Hyla gaucheri* - ESEC Grão-Pará South: CN818, CN824-827, CN836, CN864-866. *Hyla geographica* - ESEC Grão-Pará South: CN727. REBIO Maicuru: CN1550. ESEC Grão-Pará Centre: CN2094, CN2135, CN2167, CN2199, CN2208, CN2310. *Hyla leucophyllata* - ESEC Grão-Pará South: CN821, CN855. REBIO Maicuru: CN1668. *Hyla minuscula* - ESEC Grão-Pará Centre: CN2194. *Hyla minuta* - FLOTA Trombetas: CN 564-568. ESEC Grão-Pará Centre: CN2329-2339. *Hyla multifasciata* - ESEC Grão-Pará South: CN887. FLOTA Paru: CN2001. ESEC Grão-Pará Centre: CN2180, CN2316-2327. *Hyla cf. nana* - ESEC Grão-Pará South: CN684B, CN725, CN734, CN740, CN756, CN788, CN876 (all tadpoles). *Hyla punctata* - FLOTA Paru: CN1991-1994. *Hyla wavrini* - FLOTA Faro: CN113. *Osteocephalus leprieuri* - FLOTA Faro: CN82, CN84, CN206, CN307. REBIO Maicuru: CN1502, CN1533, CN1681. FLOTA Paru: CN1832. ESEC Grão-Pará Centre: CN2150, CN2368. *Osteocephalus oophagus* - FLOTA Trombetas: CN430. ESEC Grão-Pará South: CN853, CN854. ESEC Grão-Pará North: CN1269, CN1340, CN1338. *Osteocephalus taurinus* - FLOTA Faro: CN81, CN112. FLOTA Trombetas: CN431, CN513. ESEC Grão-Pará South: CN873. ESEC Grão-Pará North: CN1130. ESEC Grão-Pará Centre: CN2053, CN2077-2080,



CN2107-2109, CN2170, CN2184, CN2260, CN2286, CN2289, CN2357. *Phyllomedusa bicolor* - REBIO Maicuru: CN1591. *Phyllomedusa hypochondrialis* - REBIO Maicuru: CN1552, CN1669. ESEC Grão-Pará Centre: CN2172, CN2173, CN2312-2315. *Phyllomedusa tomopterna* - ESEC Grão-Pará Centre: CN2134. *Phyllomedusa vaillanti* - ESEC Grão-Pará North: CN1151. *Scinax boesemani* - FLOTA Faro: CN85. ESEC Grão-Pará South: CN810-817, CN822, CN823, CN856-863. ESEC Grão-Pará Centre: 2328. *Scinax garbei* - REBIO Maicuru: CN1629, CN1633, CN1654, CN1656. *Scinax nebulosus* - ESEC Grão-Pará Centre: CN2340-2348, CN2351. *Scinax proboscideus* - ESEC Grão-Pará South: CN712. *Scinax ruber* - ESEC Grão-Pará South: CN820. *Scinax* sp. n. - REBIO Maicuru: CN1628, CN1655, CN1684, CN1685, CN1689. *Scinax* gr. *ruber* - FLOTA Paru: CN1762, CN1834, CN2022.

**Leptodactylidae.** *Adenomera andreae* - FLOTA Faro: CN35, CN60, CN101, CN135, CN220, CN221, CN269, CN283, CN325. FLOTA Trombetas: CN345, CN347, CN349, CN363, CN368, CN393, CN400, CN409, CN417, CN446, CN447, CN450, CN451, CN454, CN507, CN508, CN552. ESEC Grão-Pará South: CN627. ESEC Grão-Pará North: CN1051, CN1103, CN1107, CN1165, CN1166, CN1179, CN1182, CN1203, CN1206, CN1208, CN1221, CN1235, CN1277, CN1298, CN1298, CN1308, CN1309, CN1329, CN1352, CN1354, CN1361, CN1367. REBIO Maicuru: CN1408, CN1413, CN1454, CN1463, CN1491, CN1566, CN1567, CN1609. FLOTA Paru: CN1699, CN1704-1707, CN1711, CN1712, CN1714, CN1715, CN1721, CN1722, CN1737-1741, CN1743, CN1755, CN1759, CN1764, CN1765, CN1772-1775, CN1792, CN1793, CN1800, CN1804, CN1807, CN1808, CN1812-1814, CN1816, CN1817, CN1840, CN1841, CN1847-1849, CN1853, CN1860, CN1863, CN1864, CN1872-1875, CN1882, CN1883, CN1888, CN1890, CN1894, CN1909, CN1922, CN1924, CN1946, CN1947, CN1967, CN1986, CN1987, CN2005, CN2006, CN2016-2018, CN2029, CN2033, CN2035, CN2038, CN2048. ESEC Grão-Pará

Centre: CN2064, CN2066, CN2082, CN2087, CN2124, CN2129, CN2140, CN2144, CN2145, CN2203, CN2233, CN2250, CN2258, CN2270, CN2296. *Adenomera heyeri* - ESEC Grão-Pará North: CN1226, CN1236, CN1275. *Adenomera hylaedactyla* - ESEC Grão-Pará Centre: CN2074, CN2113, CN2183, CN2196, CN2197, CN2253, CN2254, CN2291, CN2305, CN2306. *Ceratophrys cornuta* - ESEC Grão-Pará South: CN799, CN805. *Eleutherodactylus chiastonotus* - FLOTA Trombetas: CN391. REBIO Maicuru: CN1378, CN1379, CN1461, CN1556, CN1573, CN1635, CN1662, CN1670. *Eleutherodactylus fenestratus* - ESEC Grão-Pará North: CN988, CN989, CN1116, CN1117, CN1222, CN1234, CN1249, CN1299, CN1300, CN1315, CN1364. REBIO Maicuru: CN1663. *Eleutherodactylus inguinalis* - ESEC Grão-Pará North: CN1339, CN1346. REBIO Maicuru: CN1683. *Eleutherodactylus marmoratus* - ESEC Grão-Pará North: CN1075, CN1076, CN1079, CN1080, CN1089, CN1090, CN1111, CN1204, CN1265, CN1280, CN1283. *Eleutherodactylus zeuctotylus* - FLOTA Faro: CN25. FLOTA Trombetas: CN404, CN528, CN582. ESEC Grão-Pará South: CN621, CN670, CN693, CN714, CN768, CN783, CN928, CN931, CN954. ESEC Grão-Pará North: CN992, CN1000, CN1073, CN1134, CN1136, CN1242, CN1266, CN1291, CN1347, CN1350. ESEC Grão-Pará Centre: CN2103-2106, CN2234-2236, CN2371. *Leptodactylus bolivianus* - FLOTA Trombetas: CN543. ESEC Grão-Pará South: CN771, CN819, CN872. REBIO Maicuru: CN1375-1377, CN1402, CN1476, CN1480, CN1481, CN1514, CN1551, CN1554, CN1555, CN1565, CN1593-1595, CN1608, CN1627, CN1647, CN1666, CN1667, CN1682, CN1691. FLOTA Paru: CN2002-2004, 2023. *Leptodactylus knudseni* - FLOTA Faro: CN216. FLOTA Trombetas: CN392, CN401, CN428, CN432, CN472, CN509, CN532, CN540, CN561, CN569, CN595. ESEC Grão-Pará South: CN623, CN648, CN652, CN653, CN665, CN666, CN678, CN689, CN744, CN749, CN755, CN761, CN773, CN797, CN798, CN801, CN830, CN832,



CN834, CN837, CN890, CN915, CN917, CN940. ESEC Grão-Pará North: CN1223. REBIO Maicuru: CN1659, 1661. FLOTA Paru: CN1710. ESEC Grão-Pará Centre: CN2168, CN2195, CN2285, CN2353-2356. *Leptodactylus leptodactyloides* - ESEC Grão-Pará North: CN1199. *Leptodactylus longirostris* - ESEC Grão-Pará Centre: CN2072, CN2075, CN2110, CN2151, CN2174-2177, CN2209-CN2217, CN2221, CN2244, CN2251, CN2265, CN2283, CN2288, CN2290, CN2304, CN2352. *Leptodactylus myersi* - FLOTA Trombetas: CN458-471, CN514, CN536-538, CN580, CN581, CN584, CN598. ESEC Grão-Pará Centre: CN2146, CN2179, CN2245, CN2246, CN2284, CN2307, CN2369, CN2375. *Leptodactylus mystaceus* - FLOTA Faro: CN69, CN89. FLOTA Trombetas: CN437, CN529. ESEC Grão-Pará South: CN625, CN633, CN644, CN650, CN654, CN674, CN677, CN683, CN691, CN696, CN701, CN702, CN718, CN730, CN732, CN759, CN760, CN802, CN806, CN835, CN881, CN897, CN907, CN910-914, CN916, CN934, CN941, CN964. FLOTA Paru: CN1698, CN1754, CN1771, CN1795, CN1831, CN1839, CN1879, CN1880, CN1886, CN1899, CN1938, CN1940, CN1943, CN2037, CN2042. ESEC Grão-Pará Centre: CN2063. *Leptodactylus pentadactylus* - FLOTA Faro: CN57, CN59, CN114, CN120, CN137, CN139, CN143, CN163, CN194, CN235, CN326. FLOTA Trombetas: CN377, CN435, CN436, CN573. ESEC Grão-Pará South: CN920. ESEC Grão-Pará North: CN1128, CN1146-1150, CN1334, CN1337, CN1344. ESEC Grão-Pará Centre: CN2090-2093, CN2148, CN2228, CN2264, CN2364, CN2366, CN2367. *Leptodactylus petersii* - FLOTA Faro: CN90, CN118, CN208, CN211, CN215. FLOTA Trombetas: CN578. ESEC Grão-Pará South: CN762, CN886, CN896. REBIO Maicuru: CN1425, CN1596-1600, CN1632, CN1638, CN1664, CN1665, CN1688. FLOTA Paru: CN1794, CN1798, CN1799, CN1801, CN1802, CN1805, CN1850-1852, CN1896, CN1898, CN1901, CN1925, CN1926, CN1942, CN1945.

*Leptodactylus rhodomystax* Boulenger, 1884 - FLOTA Faro: CN188, CN217. ESEC Grão-Pará South: CN631, CN632, CN646, CN647, CN681, CN686-688, CN694, CN695, CN697, CN745-748, CN777, CN787, CN803, CN847, CN852, CN903, CN904, CN908, CN923, CN924, CN926, CN956-963. ESEC Grão-Pará Centre: CN2111. *Leptodactylus stenodema* - FLOTA Trombetas: CN402, CN501. *Lithodytes lineatus* - FLOTA Faro: CN311. FLOTA Trombetas: CN337, CN453, CN499, CN586. ESEC Grão-Pará South: CN763. ESEC Grão-Pará North: CN994, CN1096. REBIO Maicuru: CN1447. FLOTA Paru: CN1697, CN1719, CN1780, CN1810, CN1868, CN1908. ESEC Grão-Pará Centre: CN2068, CN2114-2119, CN2127, CN2139, CN2143, CN2252, CN2255, CN2261, CN2267, CN2274-2276, CN2293, CN2298, CN2373, CN2374. *Physalaemus ephippifer* - FLOTA Trombetas: CN338, CN429, CN484, CN503, CN562, CN597. ESEC Grão-Pará, Estanifera: CN602-610. ESEC Grão-Pará South: CN634, CN692, CN699, CN709, CN710, CN716, CN764, CN779, CN782, CN846, CN929, CN936, CN938, CN953. REBIO Maicuru: CN1384, CN1618. ESEC Grão-Pará Centre: CN2086, CN2186, CN2223, CN2259, CN2266, CN2294. **Microhylidae.** *Chiasmocleis hudsoni* - FLOTA Faro: CN42, CN80, CN210, CN212-214, CN268. ESEC Grão-Pará South: CN711. *Chiasmocleis* sp. n. - REBIO Maicuru: CN1411, CN1446, CN1484, CN1487, CN1537, CN1641, CN1642, CN1645, CN1657. FLOTA Paru: CN1767-1769, CN1869-1871, CN1881, CN1887. ESEC Grão-Pará Centre: CN2112, 2185. *Hamptophryne boliviana* - ESEC Grão-Pará South: CN796. FLOTA Paru: CN1778, CN1784-1786, CN1844, CN1845, CN2019, CN2036. *Otophryne pyburni* - ESEC Grão-Pará North: CN1358 (adult); CN1049, CN1258-1262 (tadpoles). *Synapturanus mirandaribeiroi* - FLOTA Faro: CN64, CN87, CN230, CN279. FLOTA Trombetas: CN370, CN373, CN386, CN416, CN523, CN590. ESEC Grão-Pará North: CN1025, CN1311. **Pipidae.** *Pipa snethlageae* - FLOTA Faro: CN319. *Pipa pipa* - REBIO Maicuru: CN1631,



CN1686, CN1687. FLOTA Paru: CN1855. **Ranidae.** *Rana palmipes* - ESEC Grão-Pará, Estanifera: CN600, CN601. *Gymnophiona.* **Caeciliidae.** *Caecilia tentaculata* - FLOTA Trombetas: CN355, CN372, CN375, CN381, CN485, CN500, CN574. ESEC Grão-Pará Centre: CN2138. *Microcaecilia unicolor* - FLOTA Trombetas: CN502. **Rhinatremitidae.** *Rhinatrema* sp. n.- ESEC Grão-Pará North: CN1088.

REPTILIA. Squamata. Lizards. **Gekkonidae.** *Coleodactylus amazonicus* - FLOTA Faro: CN3, CN36, CN164, CN171, CN190. FLOTA Trombetas: CN331, CN342, CN352, CN371, CN383, CN422, CN438-443, CN456, CN530, CN531, CN551, CN577, CN588, CN589. ESEC Grão-Pará South: CN629, CN630, CN660-663, CN668, CN669, CN671, CN680, CN717, CN721, CN843, CN844, CN925, CN927, CN969. ESEC Grão-Pará North: CN979, CN999, CN1037, CN1140, CN1047, CN1048, CN1172, CN1178, CN1183, CN1185, CN1194, CN1196, CN1202, CN1209, CN1246, CN1313, CN1341, CN1363. REBIO Maicuru: CN1577. FLOTA Paru: CN1915-1917, CN1923. ESEC Grão-Pará Centre: CN2069, CN2085, CN2128, CN2202, CN2205-2207, CN2249, CN2256, CN2279, CN2358. *Gonatodes annularis* - FLOTA Faro: CN129, CN254, CN298, CN327. FLOTA Trombetas: CN413, CN427. ESEC Grão-Pará North: CN998, CN1157. *Gonatodes humeralis* - FLOTA Faro: CN31, CN78, CN130, CN160, CN165, CN318. FLOTA Trombetas: CN335, CN341, CN346, CN380, CN398, CN448, CN481, CN489, CN550, CN585. ESEC Grão-Pará South: CN720, CN738, CN753, CN754, CN808. ESEC Grão-Pará North: CN1072, CN1135, CN1156, CN1184, CN1189, CN1207, CN1247, CN1327, CN1328. REBIO Maicuru: CN1423, CN1464, CN1483, CN1545, CN1546, CN1653. FLOTA Paru: CN1829, CN1833, CN1891, CN1914, CN1975. *Lepidoblepharis heyerorum* - FLOTA Faro: CN248. FLOTA Trombetas: CN351, CN387. REBIO Maicuru: CN1677. *Pseudogonatodes guianensis* - FLOTA Faro: CN5. ESEC

Grão-Pará North: CN1036, 1045, 1171, CN1302. REBIO Maicuru: CN1536, CN1610, CN1639, CN1676. *Thecadactylus rapicauda* - ESEC Grão-Pará South: CN719. ESEC Grão-Pará North: CN1248. **Gymnophthalmidae.** *Alopoglossus angulatus* - REBIO Maicuru: CN1563. FLOTA Paru: CN1865, CN1911. *Amapasaurus tetradactylus* - FLOTA Trombetas: CN332, CN356, CN362, CN369, CN388, CN408, CN424, CN445. ESEC Grão-Pará North: CN1228. *Arthrosaura kockii* - ESEC Grão-Pará North: CN1108. REBIO Maicuru: CN1383, CN1486, CN1496. *Arthrosaura reticulata* - FLOTA Faro: CN30, CN49, CN54, CN62, CN76, CN111, CN117, CN122, CN123, CN158, CN159, CN161, CN166, CN191, CN192, CN198, CN218, CN222, CN225, CN232, CN238-241, CN257, CN259, CN260, CN262, CN271-274, CN288, CN294, CN300, CN303, CN309, CN313, CN315, CN316. REBIO Maicuru: CN1395, CN1490, CN1535, CN1673. FLOTA Paru: CN1861, CN1912. *Bachia flavescens* - FLOTA Faro: CN297. FLOTA Trombetas: CN358, CN524. ESEC Grão-Pará South: CN613, CN675, CN791, CN792, CN918, CN919. ESEC Grão-Pará North: CN1102. REBIO Maicuru: CN1418, CN1451, CN1482, CN1694. FLOTA Paru: CN1718, CN1877, CN1983. ESEC Grão-Pará Centre: CN2056, CN2057, CN2125, CN2136, CN2189, CN2226, CN2271. *Bachia panoplia* - FLOTA Faro: CN167. *Cercosaura argulus* - REBIO Maicuru: CN1504. FLOTA Paru: CN1811. *Cercosaura ocellata* - ESEC Grão-Pará North: CN1318. FLOTA Paru: CN1862, CN1889, CN2028. ESEC Grão-Pará Centre: CN2071, CN2181. *Gymnophthalmus* cf. *underwoodi* - ESEC Grão-Pará Centre: CN2225. *Iphisa elegans* - ESEC Grão-Pará South: CN655, CN752, CN845, CN966. ESEC Grão-Pará North: CN1100, CN1101, CN1253, CN1254, CN1307. *Leposoma guianense* - FLOTA Faro: CN27, CN148, CN149, CN150, CN157, CN178, CN197, CN223, CN253, CN256, CN291. FLOTA Trombetas: CN344, CN407, CN423, CN444, CN452, CN493, CN592, CN599. ESEC Grão-Pará South: CN656-659, CN667, CN673, CN679, CN698, CN713, CN728,

CN731, CN741, CN772, CN778, CN780, CN786, CN851, CN900, CN905, CN930, CN955. ESEC Grão-Pará North: CN1095, CN1139, CN1159, CN1218, CN1263, CN1272, CN1296. REBIO Maicuru: CN1385, CN1386, CN1388-1391, CN1398-1401, CN1403, CN1404-1407, CN1414, CN1415, CN1420-1422, CN1428, CN1430, CN1432, CN1438, CN1439, CN1445, CN1453, CN1455, CN1457, CN1459, CN1462, CN1488, CN1489, CN1495, CN1506, CN1528, CN1542-1544, CN1547, CN1571, CN1606, CN1611, CN1612, CN1614, CN1625, CN1643, CN1644, CN1646, CN1649, CN1674, CN1675, CN1692. FLOTA Paru: CN1713, CN1981, CN2026. ESEC Grão-Pará Centre: CN2059, CN2065, CN2089, CN2227, CN2229, CN2230, CN2257, CN2268, CN2269. *Leposoma percarinatum* - FLOTA Faro: CN153, CN155, CN172-174, CN180, CN193, CN233, CN244, CN258, CN296. ESEC Grão-Pará North: CN1161, CN1252, CN1325. FLOTA Paru: CN1749, CN1770, CN1790, CN1866, CN1980, CN1985, CN1988, CN2032, CN2047. ESEC Grão-Pará Centre: CN2067, CN2084, CN2149, CN2278. *Leposoma* sp. n. - FLOTA Faro: CN147, CN151, CN152, CN154, CN179, CN209, CN249. *Neusticurus bicarinatus* - ESEC Grão-Pará North: CN1017, CN1109, CN1124, CN1238, CN1332, CN1333, CN1348, CN1369. ESEC Grão-Pará Centre: CN2231, CN2309. *Neusticurus rudis* - ESEC Grão-Pará North: CN1168, CN1284, CN1374. *Ptychoglossus brevifrontalis* - FLOTA Faro: CN124, CN224, CN263, CN245. ESEC Grão-Pará North: CN1092, CN1162. *Tretioscincus agilis* - FLOTA Faro: CN264, CN265, CN292. FLOTA Trombetas: CN389. ESEC Grão-Pará South: CN638, CN651, CN775. ESEC Grão-Pará North: CN1232. REBIO Maicuru: CN1433, CN1512, CN1538, CN1562, CN1579, CN1652, CN1672. **Polychrotidae.** *Anolis auratus* - ESEC Grão-Pará Centre: CN2073. *Anolis fuscoauratus* - FLOTA Faro: CN22, CN34, CN39, CN88, CN250, CN255, CN286. FLOTA Trombetas: CN525, CN558, CN576. ESEC Grão-Pará South: CN682, CN751, CN774, CN785, CN842, CN932. ESEC Grão-Pará North: CN1021, CN1042, CN1133. REBIO Maicuru: CN1424, CN1616, CN1660. FLOTA Paru: CN1835, CN2021. *Anolis chrysolepis* - FLOTA Faro: CN1, CN2, CN4, CN107-110, CN126-128, CN131, CN132, CN142, CN162, CN176, CN177, CN181, CN183, CN185, CN195, CN231, CN237, CN293, CN295, CN310, CN317, CN323, CN324. FLOTA Trombetas: CN406, CN494, CN547, CN591. ESEC Grão-Pará North: CN1011, CN1094, CN1104, CN1113, CN1152, CN1205, CN1230, CN1241, CN1256, CN1274, CN1282, CN1306, CN1322, CN1351, CN1362. REBIO Maicuru: CN1419, CN1431, CN1442, CN1448, CN1449, CN1468, CN1503, CN1541, CN1574, CN1576, CN1605, CN1621, CN1622. *Anolis punctatus* - ESEC Grão-Pará South: CN618. **Tropiduridae.** *Plica plica* - ESEC Grão-Pará North: CN1132. *Plica umbra* - FLOTA Faro: CN83. FLOTA Trombetas: CN374, CN382, CN497, CN498, CN505, CN544, CN545, CN560, CN572. ESEC Grão-Pará South: CN879. ESEC Grão-Pará North: CN972, CN976, CN1026, CN1030, CN1039-1041, CN1091, CN1131, CN1153, CN1193, CN1200, CN1326, CN1336, CN1349. REBIO Maicuru: CN1534, CN1619. FLOTA Paru: CN1748, CN1934. *Uranoscodon superciliosus* - FLOTA Faro: CN52. FLOTA Trombetas: CN477, CN478. ESEC Grão-Pará North: CN1190, CN1288. REBIO Maicuru: CN1601. FLOTA Paru: CN1900, CN1927. ESEC Grão-Pará Centre: CN2300. **Scincidae.** *Mabuya nigropunctata* - FLOTA Trombetas: CN473, CN526. ESEC Grão-Pará North: CN1065, CN1215, CN1216. REBIO Maicuru: CN1505. FLOTA Paru: CN1789, CN1806, CN1971, CN2007, CN2027. **Teiidae.** *Ameiva ameiva* - FLOTA Faro: CN189. ESEC Grão-Pará South: CN790. REBIO Maicuru: CN1465, CN1561, CN1651, CN1695. FLOTA Paru: CN1753, CN1776, CN1783, CN1935, CN1979, CN1982, CN2049. ESEC Grão-Pará Centre: CN2083, CN2360. *Kentropyx calcarata* - FLOTA Faro: CN219, CN247. FLOTA Trombetas: CN334, CN336, CN396, CN397, CN405, CN482, CN483, CN553. REBIO Maicuru: CN1380, CN1387, CN1392-1394, CN1409,



CN1410, CN1416, CN1417, CN1426, CN1427, CN1434-1436, CN1441, CN1456, CN1458, CN1466, CN1485, CN1507-1510, CN1513, CN1539, CN1558, CN1559, CN1564, CN1570, CN1572, CN1613, CN1615, CN1623, CN1640, CN1650, CN1678. FLOTA Paru: CN1723, CN1744-1747, CN1750-1752, CN1757, CN1781, CN1782, CN1787, CN1788, CN1791, CN1815, CN1819-1826, CN1828, CN1836-1838, CN1856, CN1857, CN1884, CN1904-1907, CN1913, CN1918-1921, CN1928, CN1929, CN1965, CN1966, CN1968-1970, CN1972, CN1974, CN1976, CN1977, CN1984, CN1989, CN1990, CN2010-2015, CN2031, CN2034, CN2039, CN2040; CN2009 (eggs). ESEC Grão-Pará Centre: CN2062. *Kentropyx striata* - ESEC Grão-Pará Centre: CN2147. Amphisbaenians. **Amphisbaenidae.** *Amphisbaena fuliginosa* - ESEC Grão-Pará South: CN612. *Amphisbaena vanzolinii* - ESEC Grão-Pará South: CN614-616, CN625, CN639, CN676, CN724, CN739, CN840, CN841, CN885, CN898, CN901, CN945-948, CN950, CN951. *Mesobaena* sp. n. - FLOTA Faro: CN7. Snakes. **Leptotyphlopidae.** *Leptotyphlops albifrons* - ESEC Grão-Pará North: CN1201. FLOTA Paru: CN1903. ESEC Grão-Pará Centre: CN2070, CN2219, CN2220. *Leptotyphlops cupinensis* - ESEC Grão-Pará South: CN767, CN902A. Typhlopidae. *Typhlops brongersmianus* - FLOTA Faro: CN282. **Boidae.** *Corallus caninus* - ESEC Grão-Pará North: CN973. *Corallus hortulanus* - FLOTA Faro: CN86, CN321. REBIO Maicuru: CN1634. *Eunectes murinus* - ESEC Grão-Pará Centre: CN2051. **Colubridae.** *Apostolepis nigrolineatus* - ESEC Grão-Pará North: CN1317. ESEC Grão-Pará Centre: CN2272. *Apostolepis quinquelineatus* - ESEC Grão-Pará South: CN902. ESEC Grão-Pará North: CN1158, CN1294. *Atractus badius* - ESEC Grão-Pará South: CN804. *Atractus torquatus* - ESEC Grão-Pará North: CN1251 CN1365, CN1368. *Chironius exoletus* - FLOTA Faro: CN320. *Chironius fuscus* - ESEC Grão-Pará North: CN1145. *Chironius multiventris cochraniae* - FLOTA Trombetas: CN556. *Chironius scurrulus* - ESEC Grão-Pará North: CN970. FLOTA Paru: CN1930. *Clelia clelia* -

FLOTA Trombetas: CN510. *Dendrophidion dendrophis* - FLOTA Faro: CN121. ESEC Grão-Pará North: CN1012, CN1023, CN1314. REBIO Maicuru: CN1532. *Dipsas catesbyi* (Sentzen, 1796) - FLOTA Trombetas: CN535. REBIO Maicuru: CN1511. *Drymoluber dichrous* - FLOTA Trombetas: CN365. FLOTA Paru: CN2025. *Erythrolamprus aesculapii* - ESEC Grão-Pará North: CN1227, CN1295. REBIO Maicuru: CN1460. *Helicops angulatus* - FLOTA Faro: CN79. FLOTA Trombetas: CN487. FLOTA Paru: CN1858, CN1859. *Imantodes cenchoa* - FLOTA Faro: CN170. FLOTA Trombetas: CN330, CN486. *Leptodeira annulata* - REBIO Maicuru: CN1382. ESEC Grão-Pará Centre: CN2361. *Liophis reginae* - ESEC Grão-Pará South: CN723, CN937. ESEC Grão-Pará North: CN993, CN1176, CN1177, CN1255, CN1366. REBIO Maicuru: CN1443, CN1493, CN1578. *Liophis typhlus* - FLOTA Trombetas: CN343, CN353, CN357, CN506, CN557, CN575, CN597. FLOTA Paru: CN1756. *Mastigodryas boddaerti* - FLOTA Trombetas: CN593. FLOTA Paru: CN1936. ESEC Grão-Pará, Estanifera: CN2052. *Oxyrhopus petola* - FLOTA Paru: CN2008. *Pseudoboa coronata* - FLOTA Faro: CN243. ESEC Grão-Pará Centre: CN2187. *Pseudoboa newwiedi* - ESEC Grão-Pará Centre: CN2218. *Pseustes poecilonotus* - ESEC Grão-Pará Centre: CN2133. *Pseustes sulphureus* - ESEC Grão-Pará South: CN878. ESEC Grão-Pará North: CN1170. REBIO Maicuru: CN1548. ESEC Grão-Pará Centre: CN2076, CN2282. *Rhinobothryum lentiginosum* - ESEC Grão-Pará Centre: CN2095. *Siphlophis compressus* - ESEC Grão-Pará Centre: CN2096. *Taeniophallus brevirostris* - FLOTA Faro: CN305. ESEC Grão-Pará North: CN977, CN1062. *Taeniophallus nicagus* - ESEC Grão-Pará North: CN1287. *Tantilla melanocephala* - FLOTA Faro: CN322. ESEC Grão-Pará South: CN933. *Thalesius viridis* - ESEC Grão-Pará North: CN1292. *Xenopholis scalaris* - REBIO Maicuru: CN1602. *Xenoxybelis argenteus* - ESEC Grão-Pará South: CN619, 635. **Elapidae.** *Micrurus averyi* - ESEC Grão-Pará North: CN1086, CN1129. *Micrurus hemprichii* - ESEC Grão-Pará Centre: CN2287. *Micrurus lemniscatus* - FLOTA Faro:



CN284. REBIO Maicuru: CN1469. *Micrurus paraensis* - ESEC Grão-Pará South: CN793. *Micrurus surinamensis* - FLOTA Faro: CN168. **Viperidae.** *Bothrops taeniatus* - FLOTA Trombetas: CN379. *Bothrops atrox* - FLOTA Trombetas: CN329, CN385, CN399, CN457, CN583. ESEC Grão-Pará North: CN971, CN1138. FLOTA Paru: CN1777, CN1830, CN1878, CN1932, CN1978, CN2020. *Lachesis muta* - REBIO Maicuru: CN1549, CN1557. FLOTA Paru: CN1809. *Chelonia*. **Testudinidae.** *Chelonoidis carbonária* - FLOTA Faro: CN328. ESEC Grão-Pará South: CN622. *Chelonoidis denticulata* - ESEC Grão-Pará South: CN664. ESEC Grão-Pará North: CN1250. REBIO Maicuru: CN1580, CN1626. **Chelidae.** *Mesoclemmys gibba* - REBIO Maicuru: CN1452. *Platemys platycephalus* - FLOTA Trombetas: CN340, CN554, CN555. **Crocodylia.** **Alligatoridae.** *Caiman crocodylus* - FLOTA Faro: CN306. *Paleosuchus trigonatus* - ESEC Grão-Pará North: CN1268. ESEC Grão-Pará Centre: CN2263.

#### NOTE ADDED IN PROOF

Franco & Ferreira (2003) mention specimens of *Thamnodynastes strigatus* (Günther, 1858) from Tapiru, Rio Paru, Municipality Almeirim (MNRJ 8735), and one from Roraima (MNRJ 0668) in the National Museum in Rio de Janeiro. As the specimens reported are far away from the known distribution of *T. strigatus*, they most likely are *T. ramonriveroi* Manzanillo & Sanchez, 2005, a species described from Venezuela, and also reported from Trinidad, Guyana, Suriname and Roraima (Brazil) by Bailey & Thomas (2007). These latter authors state that *T. ramonriveroi* is closely related to *T. strigatus* and specifically include the specimen MNRJ

0668 from Roraima in *T. ramonriveroi*. However, they do not mention the specimen from Tapiru (MNRJ 8735, which is only a head and anterior portion of body), but we think this also belongs to *T. ramonriveroi*. Bailey *et al.* (2005) described *T. lanei* from, among other places, the municipalities of Monte Alegre and Óbidos. Thus *T. ramonriveroi* and *T. lanei* can be added to the herpetofauna of CNP. Taking these new data into account, the total number of snakes known from CNP, but not collected by us, rises from 29 (see p. 48) to 31, and consequently the total number of species of reptiles known from CNP rises from 138 (see p. 71) to 140.

#### REFERENCES

BAILEY, J. R., R. A. THOMAS & N. J. SILVA, 2005. A revision of the South American snake genus *Thamnodynastes* Wagler, 1830 (Serpentes, Colubridae, Tachymenini). I. Two new species of *Thamnodynastes* from Central Brazil and adjacent areas, with a redefinition of and neotype designation for *Thamnodynastes pallidus* (Linnaeus, 1758). **Phyllomedusa** 4(2): 83-101.

BAILEY, J. R. & R. A. THOMAS, 2007 [2006]. A revision of the South American snake genus *Thamnodynastes* Wagler, 1830 (Serpentes: Colubridae, Tachymenini). II. Three new species from northern South America, with further descriptions of *Thamnodynastes gambotensis* Pérez-Santos and Moreno and *Thamnodynastes ramonriveroi* Manzanilla and Sánchez. **Memoria de la Fundacion La Salle de Ciencias Naturales** 166: 7-27.

FRANCO, F. L. & T. G. FERREIRA, 2003. Ocorência de *Thamnodynastes strigatus* (Serpentes, Colubridae) no Escudo das Guianas, Estados do Pará e Roraima, Brasil. **Phyllomedusa** 2(2): 117-119.

MANZANILLA, J. & D. SANCHEZ, 2005 [2004]. Una nueva especie de *Thamnodynastes* (Serpentes: Colubridae) del Macizo del Turimiquire, noreste de Venezuela. **Memoria de la Fundacion La Salle de Ciencias Naturales** 161-162: 61-75.

